




DOSING FOR CATS AND DOGS

Onsior™ solution for cats and dogs

1 ml/10 kg subcutaneously approximately 30 minutes before the start of surgery, and once a day for up to 2 additional days



Onsior™ solution for injection (from 2.5 kg body weight) or tablets may be used interchangeably in accordance with the indications and directions of use approved for each pharmaceutical form. Please note that the dose may be different.




Onsior™ tablets for cats (6 mg)

Musculoskeletal disorders / Orthopaedic surgery

Weight (kg)	# tablets
2.5 to < 6	●
6 to < 12	● ●

Recommended once-daily dose is 1 mg/kg with a range of 1–2.4 mg/kg body weight. Give either without food or with a small amount of food.



Onsior™ tablets for dogs

Osteoarthritis

Weight (kg)	# tablets			
	5 mg	10 mg	20 mg	40 mg
2.5 to < 5	●			
5 to < 10		●		
10 to < 20			●	
20 to < 40				●
40 to 80				● ●

Recommended dose is 1 mg/kg with a range of 1–2 mg/kg body weight, administered without food or at least 30 minutes before or after a meal.



Soft tissue surgery

Weight (kg)	# tablets			
	5 mg	10 mg	20 mg	40 mg
2.5	●			
> 2.5 to < 5		●		
5 to < 10			●	
10 to < 20				●
20 to < 40				● ●
40 to < 60				● ● ●
60 to 80				● ● ● ●

Recommended dose is 2 mg/kg with a range of 2–4 mg/kg body weight. Administer, without food, approximately 30 minutes before the operation, and once a day for up to 2 additional days.



Please refer to full product information before prescribing.



Attach double sided tape here

Onsior™ solution

- For the treatment of pain and inflammation associated with **orthopaedic or soft tissue surgery** in cats and dogs

Onsior™ tablets: CAT

- For the treatment of pain and inflammation associated with acute or chronic **musculoskeletal disorders**
- For the reduction of moderate pain and inflammation associated with **orthopaedic surgery**

Onsior™ tablets: DOG

- For the treatment of pain and inflammation associated with **soft tissue surgery and chronic osteoarthritis**

Onsior™ contains robenacoxib.

Onsior 6 mg tablets for cats are indicated for the treatment of pain and inflammation associated with acute or chronic musculoskeletal disorders in cats and for the reduction of moderate pain and inflammation associated with orthopaedic surgery in cats.

Onsior 5 mg, 10 mg, 20 mg and 40 mg tablets for dogs are indicated for the treatment of pain and inflammation associated with chronic osteoarthritis in dogs and for the treatment of pain and inflammation associated with soft tissue surgery in dogs.

Onsior 20 mg/ml solution for injection for cats and dogs is indicated for the treatment of pain and inflammation associated with orthopaedic or soft tissue surgery in dogs and for the treatment of pain and inflammation associated with orthopaedic or soft tissue surgery in cats. The dose is 2 mg/kg for dogs and cats, administered subcutaneously.

Legal Category POM V. In IE: POM.

Further information is available from the Summary of Product Characteristics.

Refer to the product packaging and leaflets for information about side effects, precautions, warnings and contraindications.

Elanco™, Onsior™ and the Diagonal Bar™ are trademarks of Elanco or its affiliates.

Use medicines responsibly www.noah.co.uk/responsible www.apha.ie

For further information contact Elanco Animal Health on +44 (0)1256 353131, or write to Elanco UK AH Limited, Form 2, Bartley Way, Bartley Wood Business Park, Hook, RG27 9XA.