Bexacat Social Toolkit

Social Media Posts



We know you are always looking for the best care for your feline patients, like prescribing Bexacat for diabetes treatment. This social toolkit contains engaging educational posts about feline diabetes and Bexacat that you can share with your clients, letting them know you have their cats' best interests at heart.

To use the posts below, follow these simple, step-by-step instructions to upload images and text. The tables below (starting on Page 2) contain the copy and image name for each post.

ATTENTION: Do not alter these posts in any way without prior approval from Elanco.

HOW TO CREATE A FACEBOOK POST: Using your phone or desktop, select "Create a post" on your clinic's Facebook page. Next, copy and paste the provided copy into the space on your new post. Then select the "Photo/video" option and attach the provided image. Be sure to add alt text by clicking the ellipsis in the upper right corner of the image. Double-check that all text tagging another account is active; this is indicated by a light blue highlight over the top of the tagged name. Copy and paste the provided copy into your new post and hit "Publish." Hashtags are not necessary on Facebook.

HOW TO CREATE AN INSTAGRAM POST: Select the "+" button at the bottom of the mobile app or select "+ Create" from the left-side navigation on your desktop to create a post on your clinic's Instagram page. Choose the selected image for the post; do not apply a filter, and proceed to the caption. To add alt text to the image, select "advanced settings" at the bottom of the screen before posting for greater accessibility. Copy and paste the provided copy into your new post and hit "Share."

To maximize character count, all recommended hashtags should be entered into the first comment of the Instagram post and **NOT** the copy. Once you have posted your image to Instagram, paste the following hashtags as a comment under your caption and hit "Post." To ensure these hashtags stay at the top of the comment section, swipe left on the comment and select the pin icon.

Recommended hashtags to paste as a comment under each post:

#FelineDiabetes #FelineDiabetesAwareness #CatDiabetes #DiabetesInCats #FelineHealth #DiabeticCatsOfInstagram #DiabeticCat

PLEASE INCLUDE THE IMPORTANT SAFETY AND LEGAL INFORMATION IN THE COPY OF ALL POSTS, INCLUDING LINKS AND TAGS. THIS INFORMATION SHOULD COME AFTER THE SUGGESTED POST COPY AND ANY INSTAGRAM HASHTAGS.

NOTE ON PRODUCT COMPLAINTS: If you receive comments on posts with a product complaint, alert Elanco by contacting us at 800-255-6826. If you do not want to monitor discussion, you can turn off comments for the provided posts.

TEXT & IMAGE ASSETS: Images can be found within the zipped folder provided for download.

Facebook Post Number	Facebook Post Copy	Facebook Post Image
Post 1: Diabetes diagnosis	Hypothetical: Your cat has just been diagnosed with diabetes. □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □	Shine a light on FELINE DIABETES Image Name: Diabetes diagnosis

may result in death. Development of these conditions should be treated promptly, including insulin administration and discontinuation of Bexacat. Do not use Bexacat in cats with diabetes mellitus who have previously been treated with insulin, who are receiving insulin, or in cats with insulin-dependent diabetes mellitus. The use of Bexacat in cats with insulin-dependent diabetes mellitus, or the withdrawal of insulin and initiation of Bexacat, is associated with an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis and death. Sudden onset of hyporexia/anorexia, lethargy, dehydration, diarrhea that is unresponsive to conventional therapy, or weight loss in cats receiving Bexacat should prompt immediate discontinuation of Bexacat and assessment for diabetic ketoacidosis, regardless of blood glucose level. Bexacat should not be initiated in cats with pancreatitis, anorexia, dehydration, or lethargy at the time of diagnosis of diabetes mellitus, as it may indicate the presence of other concurrent disease and increase the risk of diabetic ketoacidosis. Due to risk of severe adverse reactions, do not use Bexacat in cats with evidence of hepatic disease or reduced renal function. Consult a physician in case of accidental ingestion by humans.

Facebook Post Number	Facebook Post Copy	Facebook Post Image
Post 2: Diabetes treatment	A new feline diabetes diagnosis means quite a few changes for you and your cat a new routine and a few new responsibilities. In the past, this always meant twice-daily insulin injections too. But for some cats, there is a different way with Bexacat™ (bexagliflozin tablets). Now for the big question: Is your cat a "Bexacat"? This once-daily flavored tablet can make diabetes management simpler, but cats qualify only if they: ✓ Have not had insulin treatment ✓ Are healthy outside of having diabetes ✓ Are eating and drinking normally ✓ Show no evidence of significant liver or kidney disease, pancreatitis or diabetic ketoacidosis Your cat's health is our priority. If you're concerned your cat may be developing diabetes, we've got you! We'll make sure to help you and your furry friend get on an appropriate treatment plan. They might even become a "Bexacat." INDICATION: Bexacat is indicated to improve glycemic control in otherwise healthy cats with diabetes mellitus not previously treated with insulin. IMPORTANT SAFETY INFORMATION: Before using this product, it is important to read the entire product insert, including the boxed warning. See package insert for full prescribing information: https://bit.ly/3lc7ADK Cats treated with Bexacat may be at an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis, both of which may result in death. Development of these conditions should be treated promptly, including insulin administration and discontinuation of Bexacat. Do not use Bexacat in cats with diabetes mellitus who have previously been treated with insulin,	Bexact (bexagliflazin tablets) Image Name: Diabetes treatment

who are receiving insulin, or in cats with insulin-dependent diabetes mellitus. The use of Bexacat in cats with insulin-dependent diabetes mellitus, or the withdrawal of insulin and initiation of Bexacat, is associated with an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis and death. Sudden onset of hyporexia/anorexia, lethargy, dehydration, diarrhea that is unresponsive to conventional therapy, or weight loss in cats receiving Bexacat should prompt immediate discontinuation of Bexacat and assessment for diabetic ketoacidosis, regardless of blood glucose level. Bexacat should not be initiated in cats with pancreatitis, anorexia, dehydration, or lethargy at the time of diagnosis of diabetes mellitus, as it may indicate the presence of other concurrent disease and increase the risk of diabetic ketoacidosis. Due to risk of severe adverse reactions, do not use Bexacat in cats with evidence of hepatic disease or reduced renal function. Consult a physician in case of accidental ingestion by humans.

Facebook Post Number	Facebook Post Copy
Post 3: Convenience	Treating diabetic cats with insulin can be a full-time job. It's very manageable, but some may find it inconvenient. It requires up to twice-daily injections and careful storage and handling. Often, frequent changes to the dose of insulin are needed, especially right after starting treatment.
	We thought you'd be interested to know about an exciting and more convenient diabetes treatment we have at our clinic. For newly diagnosed diabetic cats not being treated with insulin who are in good health otherwise, there is Bexacat™ (bexagliflozin tablets). It's a once-daily flavored tablet that provides an alternative to traditional treatment. NO refrigeration NO needles NO dosage changes during treatment
	If you're concerned your cat may be developing diabetes, don't wait. Call us today. We have the treatments and the know-how to help keep your pet thriving. Let's chat about Bexacat!
	INDICATION: Bexacat is indicated to improve glycemic control in otherwise healthy cats with diabetes mellitus not previously treated with insulin.
	IMPORTANT SAFETY INFORMATION: Before using this product, it is important to read the entire product insert, including the boxed warning. See package insert for full prescribing information: https://bit.ly/3lc7ADK
	Cats treated with Bexacat may be at an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis, both of which may result in death. Development of these conditions should be treated promptly, including insulin administration and discontinuation of Bexacat. Do not use Bexacat in cats with

Facebook Post Image



Image Name: Convenience

diabetes mellitus who have previously been treated with insulin, who are receiving insulin, or in cats with insulin-dependent diabetes mellitus. The use of Bexacat in cats with insulin-dependent diabetes mellitus, or the withdrawal of insulin and initiation of Bexacat, is associated with an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis and death. Sudden onset of hyporexia/anorexia, lethargy, dehydration, diarrhea that is unresponsive to conventional therapy, or weight loss in cats receiving Bexacat should prompt immediate discontinuation of Bexacat and assessment for diabetic ketoacidosis, regardless of blood glucose level. Bexacat should not be initiated in cats with pancreatitis, anorexia, dehydration, or lethargy at the time of diagnosis of diabetes mellitus, as it may indicate the presence of other concurrent disease and increase the risk of diabetic ketoacidosis. Due to risk of severe adverse reactions, do not use Bexacat in cats with evidence of hepatic disease or reduced renal function. Consult a physician in case of accidental ingestion by humans.

Facebook Post Number	Facebook Post Copy	Facebook Post Image
Post 4: New option	 Bexacat™ (bexagliflozin tablets) is here. For a long time, the only way we could treat diabetic cats was injections with insulin. That's it. Now there's a needle-free option for cats who have not been treated with insulin. Bexacat is for newly diagnosed cats that are otherwise healthy, effectively controlling their blood sugar by eliminating it through the urine. Regardless of your cat's weight, the dosage is only one tablet a day.* And it's flavored! We're always thinking about how to make your and your cat's life happier, healthier and easier. That's why we are SO thrilled to offer this innovative, needle-free option. Please contact us at the clinic if you're curious to learn more about Bexacat. *Approved for cats weighing 6.6lbs. (3.0kg) or greater. INDICATION: Bexacat is indicated to improve glycemic control in otherwise healthy cats with diabetes mellitus not previously treated with insulin. IMPORTANT SAFETY INFORMATION: Before using this product, it is important to read the entire product insert, including the boxed warning. See package insert for full prescribing information: https://bit.lv/3lc7ADK Cats treated with Bexacat may be at an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis, both of which may result in death. Development of these conditions should be treated promptly, including insulin administration and discontinuation of Bexacat. Do not use Bexacat in cats with diabetes mellitus who have previously been treated with insulin, who are receiving insulin, or in cats with insulin-dependent diabetes mellitus. The use of Bexacat in cats with insulin-dependent diabetes mellitus, or the withdrawal of insulin and initiation of Bexacat, is 	Bexact beagifron tabets Image Name: New option

associated with an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis and death. Sudden onset of hyporexia/anorexia, lethargy, dehydration, diarrhea that is unresponsive to conventional therapy, or weight loss in cats receiving Bexacat should prompt immediate discontinuation of Bexacat and assessment for diabetic ketoacidosis, regardless of blood glucose level. Bexacat should not be initiated in cats with pancreatitis, anorexia, dehydration, or lethargy at the time of diagnosis of diabetes mellitus, as it may indicate the presence of other concurrent disease and increase the risk of diabetic ketoacidosis. Due to risk of severe adverse reactions, do not use Bexacat in cats with evidence of hepatic disease or reduced renal function. Consult a physician in case of accidental ingestion by humans. Bexacat, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. © 2023 Elanco or its affiliates. PM-US-23-0852

Facebook Post Number	Facebook Post Copy	Facebook Post Image
Post 5: Lifestyle	They'll still hide in boxes. They'll still chirp at birds. They'll knock over glasses, roll out the string and drag litter through the house. AND they'll still look proud while doing it. Even with diabetes, cats can be their normal selves with proper treatment. If your cat gets diagnosed with diabetes, we have your back. We'll help build a lifestyle plan for your cat that includes treatment to help manage their blood sugar. If your cat is in good health and eating and drinking normally, we can even recommend an advancement in feline medication. Bexacat™ (bexagliflozin tablets) is a needle-free, once-daily flavored tablet that provides a convenient alternative to typical insulin injections. We want all cats, even those with diabetes, to stay their frisky selves. Drop a line if you want to talk about Bexacat further. We'd love to help! INDICATION: Bexacat is indicated to improve glycemic control in otherwise healthy cats with diabetes mellitus not previously treated with insulin. IMPORTANT SAFETY INFORMATION: Before using this product, it is important to read the entire product insert, including the boxed warning. See package insert for full prescribing information: https://bit.ly/3lc7ADK Cats treated with Bexacat may be at an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis, both of which may result in death. Development of these conditions should be treated promptly, including insulin administration and discontinuation of Bexacat. Do not use Bexacat in cats with diabetes mellitus who have previously been treated with insulin, who are receiving insulin, or in cats with insulin-dependent diabetes mellitus. The use of Bexacat in cats with insulin-dependent diabetes mellitus. The use of Bexacat in cats with insulin-dependent diabetes mellitus. The use of Bexacat in cats with insulin-dependent diabetes	What does FELINE DIABETES look like? Bexact broughten tabled Image Name: Lifestyle

mellitus, or the withdrawal of insulin and initiation of Bexacat, is associated with an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis and death. Sudden onset of hyporexia/anorexia, lethargy, dehydration, diarrhea that is unresponsive to conventional therapy, or weight loss in cats receiving Bexacat should prompt immediate discontinuation of Bexacat and assessment for diabetic ketoacidosis, regardless of blood glucose level. Bexacat should not be initiated in cats with pancreatitis, anorexia, dehydration, or lethargy at the time of diagnosis of diabetes mellitus, as it may indicate the presence of other concurrent disease and increase the risk of diabetic ketoacidosis. Due to risk of severe adverse reactions, do not use Bexacat in cats with evidence of hepatic disease or reduced renal function. Consult a physician in case of accidental ingestion by humans.

Instagram Post Number	Instagram Post Copy	Instagram Post Image
Post 1: Diabetes diagnosis	Say your cat has been diagnosed with diabetes.	Shine a light on FELINE DIABETES Image Name: Diabetes diagnosis

	blood glucose level. Bexacat should not be initiated in cats with pancreatitis, anorexia, dehydration, or lethargy at the time of diagnosis of diabetes mellitus, as it may indicate the presence of other concurrent disease and increase the risk of diabetic ketoacidosis. Due to risk of severe adverse reactions, do not use Bexacat in cats with evidence of hepatic disease or reduced renal function. Consult a physician in case of accidental ingestion by humans. Bexacat, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. © 2023 Elanco or its affiliates. PM-US-23-0852	
--	--	--

Instagram Post Number	Instagram Post Copy	Instagram Post Image
Post 2: Diabetes treatment	If your cat is losing weight or drinking, urinating or eating more, these may be signs of diabetes. But don't worry! If your cat is diagnosed, we may be able to provide a convenient treatment called Bexacat™ (bexagliflozin tablets), a once-daily, flavored oral alternative to insulin injections. INDICATION: Bexacat is indicated to improve glycemic control in otherwise healthy cats with	Bexacat (bexagliflozin tablets)
	diabetes mellitus not previously treated with insulin. IMPORTANT SAFETY INFORMATION: Before using this product, it is important to read the entire product insert, including the boxed warning. See package insert for full prescribing information: @BexacatRXInfo	
	Cats treated with Bexacat may be at an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis, both of which may result in death. Development of these conditions should be treated promptly, including insulin administration and discontinuation of Bexacat. Do not use Bexacat in cats with diabetes mellitus who have previously been	
	treated with insulin, who are receiving insulin, or in cats with insulin-dependent diabetes mellitus. The use of Bexacat in cats with insulin-dependent diabetes mellitus, or the withdrawal of insulin and initiation of Bexacat, is associated with an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis and death. Sudden onset of hyporexia/anorexia, lethargy, dehydration, diarrhea that is unresponsive to conventional therapy, or weight loss in cats receiving Bexacat should prompt	Image Name: Diabetes treatment

immediate discontinuation of Bexacat and assessment for diabetic ketoacidosis, regardless of blood glucose level. Bexacat should not be initiated in cats with pancreatitis, anorexia, dehydration, or lethargy at the time of diagnosis of diabetes mellitus, as it may indicate the presence of other concurrent disease and increase the risk of diabetic ketoacidosis. Due to risk of severe adverse reactions, do not use Bexacat in cats with evidence of hepatic disease or reduced renal function. Consult a physician in case of accidental ingestion by humans. Bexacat, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. © 2023 Elanco or its affiliates. PM-US-23-0852

Instagram Post Number	Instagram Post Copy	Instagram Post Image
Post 3: Convenience	Some find treating diabetic cats with traditional treatment inconvenient. It can require twice-daily injections, refrigerating the medication and sometimes dosage adjustments. For some newly diagnosed cats, there's Bexacat TM (bexagliflozin tablets). It requires NO refrigeration, needles or dosage changes.	THE CONVENIENT ALTERNATIVE
	INDICATION: Bexacat is indicated to improve glycemic control in otherwise healthy cats with diabetes mellitus not previously treated with insulin.	
	IMPORTANT SAFETY INFORMATION: Before using this product, it is important to read the entire product insert, including the boxed warning. See package insert for full prescribing information: @BexacatRXInfo	
	Cats treated with Bexacat may be at an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis, both of which may result in death. Development of these conditions should be treated promptly, including insulin administration and discontinuation of Bexacat. Do not use Bexacat in cats with diabetes mellitus who have previously been	Bexacat Devagifican tablets
	treated with insulin, who are receiving insulin, or in cats with insulin-dependent diabetes mellitus. The use of Bexacat in cats with insulin-dependent diabetes mellitus, or the withdrawal of insulin and initiation of Bexacat, is associated with an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis and death. Sudden onset of hyporexia/anorexia, lethargy, dehydration, diarrhea that is unresponsive to conventional therapy, or weight loss in cats receiving Bexacat should prompt immediate discontinuation of Bexacat and	Image Name: Convenience

assessment for diabetic ketoacidosis, regardless of blood glucose level. Bexacat should not be initiated in cats with pancreatitis, anorexia, dehydration, or lethargy at the time of diagnosis of diabetes mellitus, as it may indicate the presence of other concurrent disease and increase the risk of diabetic ketoacidosis. Due to risk of severe adverse reactions, do not use Bexacat in cats with evidence of hepatic disease or reduced renal function. Consult a physician in case of accidental ingestion by humans. Bexacat, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. © 2023 Elanco or its affiliates. PM-US-23-0852

Instagram Post Number	Instagram Post Copy	Instagram Post Image
Post 4: New option	Peracat™ (bexagliflozin tablets) is here.	
	For a long time, insulin was the only option for diabetic cats. We are lucky to finally have an alternative that effectively controls blood sugar without needles. It's one flavored tablet, once a day! Want to learn more? Please call us to make an appointment.	
	INDICATION: Bexacat is indicated to improve glycemic control in otherwise healthy cats with diabetes mellitus not previously treated with insulin.	
	IMPORTANT SAFETY INFORMATION: Before using this product, it is important to read the entire product insert, including the boxed warning. See package insert for full prescribing information: @BexacatRXInfo	
	Cats treated with Bexacat may be at an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis, both of which may result in death. Development of these conditions should be treated promptly, including insulin administration and	
	discontinuation of Bexacat. Do not use Bexacat in cats with diabetes mellitus who have previously been treated with insulin, who are receiving insulin, or in cats with insulin-dependent diabetes mellitus. The	Bexacat (bexaguiflozin tablets) Image Name: New option
	use of Bexacat in cats with insulin-dependent diabetes mellitus, or the withdrawal of insulin and initiation of Bexacat, is associated with an increased risk of diabetic ketoacidosis or euglycemic diabetic	
	ketoacidosis and death. Sudden onset of hyporexia/anorexia, lethargy, dehydration, diarrhea that is unresponsive to conventional therapy, or weight loss in cats receiving Bexacat should prompt	

assessment for diabetic ketoacidosis, regardless of blood glucose level. Bexacat should not be initiated in cats with pancreatitis, anorexia, dehydration, or lethargy at the time of diagnosis of diabetes mellitus, as it may indicate the presence of other concurrent disease and increase the risk of diabetic ketoacidosis. Due to risk of severe adverse reactions, do not use Bexacat in cats with evidence of hepatic disease or reduced renal function. Consult a physician in case of accidental ingestion by humans. Bexacat, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. © 2023 Elanco or its affiliates. PM-US-23-0852

Instagram Post Number	Instagram Post Copy	Instagram Post Image
Post 5: Lifestyle	Watching cats doing cat things is pure joy. We Even diabetic cats can stay their frisky selves with proper treatment. If your cat gets diagnosed with diabetes, we can recommend Bexacat™ (bexagliflozin tablets), a once-daily, flavored tablet that's an alternative to traditional treatment. INDICATION: Bexacat is indicated to improve glycemic control in otherwise healthy cats with diabetes mellitus not previously treated with insulin. IMPORTANT SAFETY INFORMATION: Before using this product, it is important to read the entire product insert, including the boxed warning. See package insert for full prescribing information: □BexacatRXInfo Cats treated with Bexacat may be at an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis, both of which may result in death. Development of these conditions should be treated promptly, including insulin administration and discontinuation of Bexacat. Do not use Bexacat in cats with diabetes mellitus who have previously been treated with insulin, who are receiving insulin, or in cats with insulin-dependent diabetes mellitus. The use of Bexacat in cats with insulin-dependent diabetes mellitus, or the withdrawal of insulin and initiation of Bexacat, is associated with an increased risk of diabetic ketoacidosis or euglycemic diabetic ketoacidosis and death. Sudden onset of hyporexia/anorexia, lethargy, dehydration, diarrhea that is unresponsive to conventional therapy, or weight loss in cats receiving Bexacat should prompt immediate discontinuation of Bexacat and	What does FELINE DIABETES look like? Bexacat bacylitan talan Image Name: Lifestyle

assessment for diabetic ketoacidosis, regardless of blood glucose level. Bexacat should not be initiated in cats with pancreatitis, anorexia, dehydration, or lethargy at the time of diagnosis of diabetes mellitus, as it may indicate the presence of other concurrent disease and increase the risk of diabetic ketoacidosis.

Due to risk of severe adverse reactions, do not use
Bexacat in cats with evidence of hepatic disease or reduced renal function. Consult a physician in case of accidental ingestion by humans.

Bexacat, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. © 2023 Elanco or its affiliates. PM-US-23-0852