



Elanco™


FLUBENVET.™
GET WORMS SORTED
AT EVERY LIFE STAGE.

Take control of your bird's health, welfare and productivity with the **only** dewormer that targets **all life stages** – not just the adult stage.¹

Elanco

Flubenvet™

Only **Flubenvet** kills all life stages of **all** the major worm species, including **larvae and eggs**¹



Worms that have reached **adulthood** live **inside** the bird, **reproduce** and can **cause** a variety of symptoms, including:

- **Poor body weight and enteritis**
- **Poor hatchability**
- **Anaemia**
- **Reduced laying rates**
- **Reduced egg weights**
- **Respiratory infections (gapes)**

Worm burdens can cause the death of up to 19% of released pheasants in the UK²

Why is **broad-spectrum** deworming with Flubenvet important?

Parasitic worms can damage the health, welfare and performance of your birds. Controlling these parasites is a problem you face every season. Flubenvet controls all of the key species of worms affecting your birds.

Gapeworms – A major cause of death and respiratory infections in both pheasants and partridges. These worms inhabit the lungs and windpipe, causing birds to 'gape' or 'snick' as they struggle to breathe.

Roundworms – Live in the small intestine and can be up to 2 inches long. In heavy infestations, these worms may be visible in faeces.

Hairworms – Barely visible to the naked eye, but can cause significant commercial damage, even in moderate infestations.

Caecal worms – Smaller worms inhabiting the lower end of the gut. Infected birds often appear asymptomatic.

Adult worms are only part of the problem

The immature life stages of parasitic worms also contribute to the disease. Gapeworm larvae, for example, cause damage whilst burrowing from the intestines, through the lungs towards the windpipe. Infectious larvae can survive in earthworms for up to three years, providing a continuous pool of reinfection across multiple seasons, unless birds are effectively dewormed and ground is rested appropriately to help reduce the burden.

Targeting every stage of the worm lifecycle is crucial when trying to break the rapid worm lifecycle.

Flubenvet is administered conveniently in feed for 7 days

When weather conditions are wet, then it's likely that your birds will seek water from puddles instead of drinkers. Therefore, using Flubenvet in feed helps you to be assured that your birds are receiving the correct dose of dewormer.



Does your wormer target **ALL** worms at **ALL** life stages?

	Target poultry species	Gapeworm (<i>Syngamus trachea</i>)	Roundworm (<i>Ascaridia galli</i>)	Hairworm (<i>Capillaria spp.</i>)	Caecal worm (<i>Heterakis gallinarum</i>)	Caecal threadworm (<i>Trichostrongylus tenuis</i>)
FLUBENVET PREMIX (Flubendazole in feed)	Pheasants, Partridges, Geese, Turkeys, Chickens	✓	✓	✓	✓	✓
FLUBENMIX (Flubendazole in feed)	Pheasants, Partridges, Geese, Turkeys, Chickens	Adult stages only	Adult stages only	Adult stages only	Adult stages only	Adult stages only
GALLIFEN PREMIX (Fenbendazole in feed)	Pheasants, Chickens	✗	Adult stages only in CHICKENS	✗	Adult + L5 stages only in CHICKENS Adult stages only in PHEASANTS	✗
GALLIFEN (Fenbendazole suspension)	Pheasants, Chickens	✗	Adult stages only in CHICKENS	Adult stages only in CHICKENS	Adult stages only	✗



Scan to discover more about Flubenvet

1. Flubenvet 5% SPC. 2. Madden J. et al (2018), Why do pheasants released in the UK die and how can we best reduce their natural mortality? European Journal of Wildlife Research. Available at: <https://doi.org/10.1007/s10344-018-1199-5>.

Flubenvet 5% w/w Premix for Medicated Feeding Stuff contains 50 mg/g flubendazole. Legal Category POM-VPS. Birds must not be slaughtered for human consumption during treatment. Chickens, turkeys, geese, partridges and pheasants: Withdrawal period: Meat: 7 days Chickens eggs: zero days. To be supplied only on prescription.

Advice on the use of Flubenvet or alternative medicines must be sought from the medicine prescriber. **Use medicines responsibly** www.noah.co.uk/responsible. Elanco UK AH Limited, Form 2, Bartley Way, Bartley Wood Business Park, Hook RG27 9XA. Telephone: 01256 353131. Email: elancouk@elanco.com. Flubenvet, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates. Other product names are trademarks of their respective owners. © 2023 Elanco or its affiliates. Date of preparation: 03/23. PM-UK-23-0094

Elanco

Flubenvet™