

Elanco

Optaflexx®

AT-A-GLANCE

PRODUCT SUMMARY

Optaflexx® is a feed ingredient that can increase live weight gain 22 lbs and hot carcass weight 20 lbs when fed to steers at 300 mg/hd/d for the last 28 to 42 days of finishing. Optaflexx is a high-reward technology that adds pounds of beef while maintaining an animal's ability to grade Choice or better.¹



Optaflexx puts more money in your pocket by adding pounds of beef while providing you the freedom to make timely decisions managing and marketing cattle as the market fluctuates.

DOSE FLEXIBILITY

Optaflexx allows feeders to select the dose level that meets their performance objectives and easily respond to market signals and the Choice-Select spread.

ENHANCED ROI

Optaflexx enhances ROI whether selling live cattle, carcasses or carcasses on a value-based grid.¹

REWARD WITHOUT RESTRAINT

A zero-day withdrawal facilitates ease of implementation and allows flexibility in end-point management. Optaflexx is also accepted by all major packers.

OPTAFLEXX GUIDELINES & LABEL DIRECTIONS

The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

Optaflexx: Complete feed

Caution: Not for animals intended for breeding.

For increased rate of weight gain and improved feed efficiency in cattle fed in confinement for slaughter: Feed 8.2 to 24.6 g/ton of ractopamine hydrochloride (90% DM basis) continuously in a complete feed to provide 70 to 430 mg/hd/d for the last 28 to 42 days on feed.

For increased rate of weight gain, improved feed efficiency and increased carcass leanness in cattle fed in confinement for slaughter: Feed 9.8 to 24.6 g/ton of ractopamine hydrochloride (90% DM basis) continuously in a complete feed to provide 90 to 430 mg/hd/d for the last 28 to 42 days on feed.

Optaflexx: Top dress

For increased rate of weight gain and improved feed efficiency in cattle fed in confinement for slaughter: Feed 70 to 400 mg/hd/d of ractopamine hydrochloride (90% DM basis) continuously in a minimum of 1.0 lb/hd/d top dress Type C medicated feed (maximum 800 g/ton ractopamine hydrochloride) during the last 28 to 42 days on feed.

¹ Elanco Animal Health. Data on file.

Elanco