



Product Name: COBALIFE VB12 PLUS SELENIUM FOR SHEEP AND CATTLE  
APVMA Approval No: 61465 / 130880

Label Name:	COBALIFE VB12 PLUS SELENIUM FOR SHEEP AND CATTLE
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY
Constituent Statements:	4 mg/mL Selenium as Sodium Selenate 2 mg/mL HYDROXOCOBALAMIN (present as HYDROXOCOBALAMIN ACETATE Vitamin B12)
Claims:	For the treatment and control of Vitamin B12 and selenium deficiency in sheep and cattle
Net Contents:	200 mL 500 mL
Directions for Use:	
Restraints:	
Contraindications:	This product is contraindicated for use when selenium intake from pasture is high or selenium is provided by other means (such as top dressing, vaccine, pellets or selenium drenches) if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian.
Precautions:	This product should not be administered to sheep with known liver damage concurrently grazing plants which contain pyrrolizidine alkaloids, such as Patterson's Curse or heliotrope.
Side Effects:	
Dosage and Administration:	Use the contents within 24 hours of first broaching. Discard the unused portion. Give dose by subcutaneous injection only.

	<p>LAMBS 0.5 mL at marking or weaning  SHEEP 1 - 1.5 mL pre-lambing  CALVES 2 – 3 mL from 2 months of age.  COWS 4 – 6 mL pre-calving</p> <p>The frequency of dosing should be related to the severity of the deficiencies. For instance, for sheep in extremely deficient areas, repeat dose every 2-3 months. In severely selenium deficient areas, use a calculated dose of 0.1 mg Se/kg body weight. In less deficient areas, repeat dose every 5-6 months or as found necessary. Consult your animal health adviser for advice specific to your flock or herd.</p> <p>CAUTION: Avoid carcass damage</p> <ul style="list-style-type: none"> <li>• Sterilise all injection apparatus by boiling (or equivalent) before use. Avoid use of strong disinfectants on apparatus.</li> <li>• Maintain cleanliness at all times.</li> <li>• Keep needles sharp and clean. Replace frequently.</li> <li>• Use shortest needles possible, certainly not exceeding 15 mm.</li> <li>• As far as possible, avoid injection of animals during wet weather or under dusty conditions.</li> <li>• This product should be injected only under the skin.</li> <li>• If possible, inject high on the neck behind the ear.</li> </ul>
--	---

General Directions:	
---------------------	--

Withholding Periods:	MEAT: Zero (0) days MILK: Zero (0) days
----------------------	--

Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days. Before using this product, confirm the current ESI from Elanco Australasia Pty Ltd on 1800 995 709 or the APVMA website ( <a href="http://www.apvma.gov.au/residues">www.apvma.gov.au/residues</a> ).
---------------	---

Safety Directions:	
--------------------	--

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.
-------------------------	---

First Aid Warnings:	
---------------------	--

Additional User Safety:	
-------------------------	--

Environmental Statements:	
---------------------------	--

Disposal:	Dispose of container by wrapping with paper and putting in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' container.
-----------	--

Storage:

Store below 25°C (air conditioning). Protect from light.  
Store opened product below 25°C