

Product Name: QUICKBAYT SPRAY FLY BAIT
APVMA Approval No: 61737/119534



Label Name:	QUICKBAYT SPRAY FLY BAIT
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	100 g/kg IMIDACLOPRID 0.84 g/kg (Z)-9 TRICOSENE
Mode of Action:	GROUP 4A INSECTICIDE
Statement of Claims:	A water dispersible granule formulation to be mixed with water and used as a spray bait formulation for the control of adult houseflies and lesser houseflies (including organophosphate resistant strains) in commercial, industrial and domestic areas, as per the DIRECTIONS FOR USE. Will visibly kill flies for 4 weeks.
Net Contents:	Pack size range 12.5 g - 1.5 kg Outer packs 250 g: contains 4 x 62.5 g measure packs which it is illegal to sell separately. 1 kg: contains 4 x 250 g measure packs which it is illegal to sell separately. Measure packs 62.5 g and 250 g Not to be sold separately. Before use read all directions on the outer pack.
Restrains:	DO NOT apply in a manner which may cause contamination of animal feed DO NOT apply spray treatment outdoors if rain is expected within 48 hours DO NOT apply onto material which birds are likely to feed on directly (i.e. edible refuse)
Directions for Use:	This section contains file attachment.

--	--

Other Limitations:	
--------------------	--

Withholding Periods:	
----------------------	--

Trade Advice:	
---------------	--

General Instructions:	<p>QuickBayt Spray can be applied in many locations. Avoid cool or windy locations because flies prefer warm sites for resting. For best results time applications to begin at the start of the season before fly populations have reached their peak.</p> <p>Fly control should be considered part of an integrated strategy which incorporates good sanitation, appropriate management of breeding sites (eg. manure and other decaying organic matter) and exclusion measures for preventing access of flies into internal areas.</p> <p>Note on Application Avoid applying this product to dusty/dirty surfaces. If the treated area becomes covered by dust/dirt, the product can be reactivated by gently spraying water mist on the treated surfaces. DO NOT store diluted insecticide. Prepare a fresh solution as necessary. DO NOT spray the moving parts of any machinery, electric motors or switchgear with the water-based product.</p> <p>Re-baiting Re-apply bait when flies are no longer visibly affected in areas where the product has been applied (usually after about 4 to 6 weeks). For season-long control, monthly applications may be required, depending on the situation and duration of the fly season.</p> <p>Cleaning Spray Equipment Clean spray equipment by rinsing thoroughly with lukewarm water. DO NOT allow washings to enter sewer drains or nearby water bodies.</p> <p>WARNING: In some situations mould may develop (some time after application) on areas where this product has been used. If appearance of mould on a wall or fixture is a concern in the area of application then it is recommended that a surrogate target such as a disposable piece of timber, card or other surface is treated in its place. A suitable detergent and cloth may also be used to remove mould in some areas. Dispose of cloth after use.</p>
-----------------------	--

Resistance Warning:	<p>For insecticide resistance management, QuickBayt Spray is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to QuickBayt Spray and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if QuickBayt Spray or other Group 4A insecticides are used repeatedly. The effectiveness of QuickBayt Spray on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of QuickBayt Spray to control resistant insects. For further information on managing insecticide resistance contact your local supplier or Bayer CropScience Pty Ltd Environmental Science representative.</p>
---------------------	---

Precautions:	<p>DO NOT allow contact with raw or processed food, utensils or packaging. DO NOT apply to surfaces within the reach of animals and children.</p>
--------------	--

	DO NOT spray directly on animals, feed or drinking supplies, or on milking equipment. DO NOT apply past the point of run-off.
Protections:	<p>PROTECTION OF LIVESTOCK DO NOT apply bait in locations accessible to domestic animals or livestock. Imidacloprid is dangerous to bees. QuickBayt Spray will kill bees in hives which are over-sprayed or reached with spray drift.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND THE ENVIRONMENT DO NOT apply bait in locations accessible to birds. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Clean up spillage promptly.</p>
Storage and Disposal:	<p>Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight.</p> <p>For sachets Single-rinse or shake remainder into spray tank/water. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on site. Do not burn empty containers or product.</p> <p>For plastic bottles Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. Do not reuse empty container for any other purpose.</p>
Safety Directions:	When opening the container, preparing and using the prepared spray or paste wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat and elbow-length chemical resistant gloves. Wash hands after use.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre Phone Australia 13 11 26.
First Aid Warnings:	

DIRECTIONS FOR USE**Restrictions**

DO NOT apply in a manner which may cause contamination of animal feed

DO NOT apply spray treatment outdoors if rain is expected within 48 hours

DO NOT apply onto material which birds are likely to feed on directly (i.e. edible refuse)

SITUATION	PEST	RATE	CRITICAL COMMENTS
This product is not a conventional residual spray and therefore does not work by flies simply contacting a sprayed surface. The attractants within this bait mean that flies will be attracted to and subsequently attempt to feed on treated surfaces; they will then pick up a lethal dose. Because of this bait function, only portions of vertical surfaces need to be treated.			
<p>Outside</p> <p>Abattoirs, animal housing, domestic dwellings, garbage containers, food processing plants and other industrial and commercial areas</p> <p>Around</p> <p>Industrial sites, feedlot manure heaps, garbage dumps, poultry farms and animal housing (ONLY in areas inaccessible to livestock, domestic animals and children)</p> <p>Inside</p> <p>Commercial and industrial facilities and animal houses (ONLY in areas inaccessible to livestock, domestic animals and children)</p>	<p>Nuisance flies including:</p> <p>Adult houseflies (<i>Musca domestica</i>) and Lesser houseflies (<i>Fannia canicularis</i>)</p>	<p>12.5 g/100 mL water</p> <p>(spray on application)</p>	<p>Mix 12.5 g of QuickBayt Spray with 100 mL of water and apply as a coarse spray to areas where flies rest. Maintain agitation of suspension during application to ensure thorough mixing.</p> <p>For floor areas of 5 m² it is recommended that 100mL of mix is applied to approximately 1/3 of the surrounding (mainly vertical) surfaces. Any floor areas greater than 5m² needs to be recalculated on the basis of 100mL mixed bait per 5 m² floor area.</p> <p>Note: Around feedlot manure heaps: DO NOT apply bait on manure heaps or dispose of treated targets with manure.</p> <p>See also Note on Application below.</p>
		<p>12.5 g/10 mL water</p> <p>(paint on application)</p>	<p>To prepare a paste for painting, mix 12.5 g of QuickBayt Spray with 10 mL of water. The mix can be applied with a brush to spots where flies rest (eg. walls, joists, window ledges, stable fixtures). Additionally it can be painted onto pieces of timber, or sheets of cardboard or fabric which are hung in areas frequented by flies. After use the brush can be cleaned with warm water.</p> <p>For floor areas of 5 m² it is recommended that 10 mL of mix is applied in 3 to 5 stripes to the surrounding (mainly vertical) surfaces. Any areas greater than 5 m² need to be recalculated on the basis of 10 mL mixed bait per 5 m² floor area.</p> <p>See also Note on Application below.</p>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

DIRECTIONS FOR USE**Restraints**

DO NOT apply in a manner which may cause contamination of animal feed

DO NOT apply spray treatment outdoors if rain is expected within 48 hours

DO NOT apply onto material which birds are likely to feed on directly (i.e. edible refuse)

SITUATION	PEST	RATE	CRITICAL COMMENTS
This product is not a conventional residual spray and therefore does not work by flies simply contacting a sprayed surface. The attractants within this bait mean that flies will be attracted to and subsequently attempt to feed on treated surfaces; they will then pick up a lethal dose. Because of this bait function, only portions of vertical surfaces need to be treated.			
<p>Outside</p> <p>Abattoirs, animal housing, domestic dwellings, garbage containers, food processing plants and other industrial and commercial areas</p> <p>Around</p> <p>Industrial sites, feedlot manure heaps, garbage dumps, poultry farms and animal housing (ONLY in areas inaccessible to livestock, domestic animals and children)</p> <p>Inside</p> <p>Commercial and industrial facilities and animal houses (ONLY in areas inaccessible to livestock, domestic animals and children)</p>	<p>Nuisance flies including:</p> <p>Adult houseflies (<i>Musca domestica</i>) and Lesser houseflies (<i>Fannia canicularis</i>)</p>	<p>62.5 g/500 mL water</p> <p>(spray on application)</p>	<p>Mix 62.5 g of QuickBayt Spray with 500mL of water and apply as a course spray to areas where flies rest. Maintain agitation of suspension during application to ensure thorough mixing.</p> <p>For floor areas of 25 m² it is recommended that 500mL of mix is applied to approximately 1/3 of the surrounding (mainly vertical) surfaces. Any floor area greater than 25 m² needs to be recalculated on the basis of 500mL mixed bait per 25 m² floor area.</p> <p>Note: Around feedlot manure heaps: DO NOT apply bait on manure heaps or dispose of treated targets with manure.</p> <p>See also Note on Application below.</p>
		<p>62.5 g/50 mL water</p> <p>(paint on application)</p>	<p>To prepare a paste for painting, mix 62.5 g of QuickBayt Spray with 50 mL of water. The mix can be applied with a brush to spots where flies rest (eg. walls, joists, window ledges, stable fixtures). Additionally it can be painted onto pieces of timber, or sheets of cardboard or fabric which are hung in areas frequented by flies. After use the brush can be cleaned with warm water.</p> <p>For floor areas of 25 m² it is recommended that 50 mL of mix is applied in 12 to 25 stripes to the surrounding (mainly vertical) surfaces. Any areas greater than 25 m² need to be recalculated on the basis of 50 mL mixed bait per 25 m² floor area.</p> <p>See also Note on Application below.</p>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

QuickBayt® Spray Fly Bait**Directions for Use****250 g pack**

(for outer packs containing 250 g measure packs)

DIRECTIONS FOR USE (for outer packs containing 250 g measure packs)**Restraints**

DO NOT apply in a manner which may cause contamination of animal feed

DO NOT apply spray treatment outdoors if rain is expected within 48 hours

DO NOT apply onto material which birds are likely to feed on directly (i.e. edible refuse)

SITUATION	PEST	RATE	CRITICAL COMMENTS
This product is not a conventional residual spray and therefore does not work by flies simply contacting a sprayed surface. The attractants within this bait mean that flies will be attracted to and subsequently attempt to feed on treated surfaces; they will then pick up a lethal dose. Because of this bait function, only portions of vertical surfaces need to be treated.			
<p>Outside</p> <p>Abattoirs, animal housing, domestic dwellings, garbage containers, food processing plants and other industrial and commercial areas</p> <p>Around</p> <p>Industrial sites, feedlot manure heaps, garbage dumps, poultry farms and animal housing (ONLY in areas inaccessible to livestock, domestic animals and children)</p> <p>Inside</p> <p>Commercial and industrial facilities and animal houses (ONLY in areas inaccessible to livestock, domestic animals and children)</p>	<p>Nuisance flies including:</p> <p>Adult houseflies (<i>Musca domestica</i>) and Lesser houseflies (<i>Fannia canicularis</i>)</p>	<p>250 g/2 L water</p> <p>(spray on application)</p>	<p>Mix 250 g of QuickBayt Spray with 2 L of water and apply as a course spray to areas where flies rest. Maintain agitation of suspension during application to ensure thorough mixing.</p> <p>For floor areas of 100 m² it is recommended that 2 L of mix is applied to approximately 1/3 of the surrounding (mainly vertical) surfaces. Any floor area greater than 100m² needs to be recalculated on the basis of 2 L mixed bait per 100 m² floor area.</p> <p>Note: Around feedlot manure heaps: DO NOT apply bait on manure heaps or dispose of treated targets with manure.</p> <p>See also Note on Application below.</p>
		<p>250 g/200mL water</p> <p>(paint on application)</p>	<p>To prepare a paste for painting, mix 250 g of QuickBayt Spray with 200 mL of water. The mix can be applied with a brush to spots where flies rest (eg. walls, joists, window ledges, stable fixtures). Additionally it can be painted onto pieces of timber, or sheets of cardboard or fabric which are hung in areas frequented by flies. After use the brush can be cleaned with warm water.</p> <p>For floor areas of 100 m² it is recommended that 200 mL of mix is applied in 50 to 100 stripes to the surrounding (mainly vertical) surfaces. Any areas greater than 100m² need to be recalculated on the basis of 200 mL mixed bait per 100 m² floor area.</p> <p>See also Note on Application below.</p>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**