

ACTIVE CONSTITUENTS:

Each litre contains: Calcium Chloride Dihydrate 283.1g, Magnesium Chloride Hexahydrate 130.8g and Propylene Glycol 307.7g.

Each mL contains: Calcium 77.2mg, Magnesium 15.6mg and Propylene Glycol 307.7mg.

READ ENTIRE LABEL BEFORE USE. May be harmful if swallowed.

HANDLING PRECAUTIONS:

Wash hands and exposed skin before meals and after work.

WITHOLDING PERIODS: Milk and Meat: NIL

STORAGE: Store below 30°C (room temperature). Protect from light.

DISPOSAL: Preferably dispose of the product by use. Otherwise dispose of product and packaging in an approved landfill or other approved facility.

Registered pursuant to the ACVM Act 1997, No A9642

See www.foodsafety.govt.nz for registration conditions.

Elanco New Zealand

106 Wiri Station Road, Manukau, Auckland 2104, New Zealand

Customer Info Line: 0800 446 121


WARNING

KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

ORAL-MAX™**Oral calcium supplement
and anti-ketogenic agent**

To aid in the prevention and treatment of
milk fever and ketosis in cattle. Also to supplement
dietary calcium when a deficiency may occur.

Each 650mL bottle contains:
Calcium 50.2g, Magnesium 10.1g
and Propylene Glycol 200g.



DIRECTIONS FOR USE: *By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.*

FOR ORAL USE ONLY

Administer as a drench or mix in the drinking water or feed.

Take care when drenching, particularly if the cow exhibits any sign of paralysis. Direct the neck of the bottle over the tongue towards the back of the throat and gradually squeeze the contents from the bottle to assist swallowing. Avoid accidental administration into the windpipe and lungs.

Milk Fever – Prevention Administer one bottle (650mL) 24 hours before calving, one bottle at calving, one bottle 10 – 12 hours after calving and one bottle 24 hours after calving.

Milk Fever – Treatment In conjunction with intravenous calcium treatment: Administer one bottle (650mL) as soon as the cow is able to stand and can swallow normally. Repeat the same dosage (650mL) 10 – 12 hours later.

Ketosis Administer one bottle (650mL) twice daily.

General Calcium Supplementation Give one bottle (650mL) morning and night for two days.

Note: Avoid giving extra calcium daily to pregnant cows in the transition period (4 weeks before parturition) as this may predispose to milk fever.

BATCH:

EXPIRY:

650mL

90198318

85mm deep

245mm wide

PRODUCT INFO

BLUE #:	700437APB	Product Name:	ORAL MAX
Item Code:	90198318	Component:	Label
Product Code:	N/A	Pack Size:	650mL
Previous Item Code	81490635		

ARTWORK INFO

Template:	245(W) x 85(D)mm	Packaging Spec(s):	N/A
Barcodes/Type:	EAN13: 9415530006254	Add. Info:	N/A
		Minimum Core Data Point Size:	6pt
GTIN:	GTIN Not Required	Proof #	P4a By/Date
			CY 08-Nov-2021

ELANCO ARTWORK LEGEND v13