50 Doses 100 mL

Clostridium Perfringens Type C-Escherichia Coli Bacterin-Toxoid



For Use In Animals Only

INDICATIONS: This product has been shown to be effective for the vaccination of healthy pregnant swine to provide passive immunity in piglets against disease caused by *Clostridium perfringens* Type C and K88, K99, 987P, and F41 piliated *Escherichia coli*. The duration of immunity for these antigens has not been established. For more information go to *productdata.aphis.usda.gov*.

ADJUVANT: Aluminum hydroxide.

DIRECTIONS: Shake well before using. Administer 2 mL intramuscularly at 5 and 2 weeks prior to farrowing. It is essential that newborn pigs receive colostrum from the vaccinated dam. Revaccinate prior to each subsequent farrowing.

PRECAUTIONS: Store out of direct sunlight at 2° to 8°C (35° to 46°F). DO NOT FREEZE. Do not mix with other products, except as specified on this label. Use entire contents when first opened. In case of human exposure, contact a physician. Do not vaccinate within 21 days prior to slaughter. Anaphylactic reactions may occur. Symptomatic treatment: Epinephrine.

VLN/PCN: 196/7890.00 Elanco US Inc., Fort Dodge, IA 50501 USA; 1-800-428-4441

Pili Shield, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.