



Product Name: BAYMEC LV POUR-ON BROAD SPECTRUM ENDECTOCIDE FOR DAIRY AND BEEF CATTLE
 APVMA Approval No: 64610/135073

Label Name:	BAYMEC LV POUR-ON BROAD SPECTRUM ENDECTOCIDE FOR DAIRY AND BEEF CATTLE
-------------	--

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	10 mg/mL IVERMECTIN Solvent: 150 g/L Diethylene glycol monobutyl ether
-------------------------	---

Claims:	<p>For the treatment and control of ivermectin sensitive internal and external parasites of beef and dairy cattle. BAYMEC LV Pour-On contains ivermectin, which has a broad spectrum of activity against most internal and external parasites with sensitivity to ivermectin. Ivermectin is a member of the macrocyclic lactones (anthelmintic family of chemicals).</p> <p>Indications For the treatment and control of adult and some immature stages of:</p> <p>Gastro-intestinal roundworms: Barbers Pole Worm (<i>Haemonchus placei</i>) Hookworm (<i>Bunostomum phlebotomum</i>) Intestinal Threadworm (<i>Strongyloides papillosus</i>) Large Bowel Worm (<i>Oesophagostomum radiatum</i>) Small Intestinal Worm (<i>Cooperia oncophora</i>) (<i>Cooperia punctata</i>) Small Brown Stomach Worm (<i>Ostertagia ostertagi</i>) (including inhibited stages) Stomach Hair Worm (<i>Trichostrongylus axei</i>) Thin Necked Intestinal Worm (<i>Nematodirus spathiger</i>) Whipworm (<i>Trichuris</i> spp.) Lungworms (<i>Dictyocaulus viviparus</i>) Sucking and Biting lice Biting lice (<i>Bovicola bovis</i>) Little sucking lice (<i>Solenopotes capillatus</i>) Long nose sucking lice (<i>Linognathus vituli</i>) Short nosed sucking lice (<i>Haematopinus eurysternus</i>) Mites (<i>Chorioptes bovis</i>, <i>Sarcoptes scabiei</i> var. <i>bovis</i>) Buffalo Fly (<i>Haematobia irritans exigua</i>) Cattle Tick (<i>Rhipicephalus (Boophilus) australis</i>) Maximum tick efficacy is achieved 4-5 days after treatment. In planning a strategic tick control program, it should be noted that viable eggs may have been laid by engorged female ticks that drop from cattle in the initial days after treatment. BAYMEC LV Pour-On</p>
---------	--

	<p>should be alternated with conventional dips or Pour-On's when the program requires repeat treatments.</p> <p>Sustained Activity</p> <p>BAYMEC LV Pour-On applied at the recommended dose of 1 mL per 20 kg bodyweight effectively controls infection with <i>Ostertagia</i> spp. (Small brown stomach worm) acquired up to 14 days after treatment, with <i>Dictyocaulus viviparus</i> (Lungworm) acquired up to 28 days after treatment and <i>Haematobia irritans exigua</i> (Buffalo fly) up to 28 days after treatment. Resistance may develop to any chemical.</p>
--	--

Net Contents:	<p>10L</p> <p>12.5L</p> <p>15L</p> <p>2.5L</p> <p>20L</p> <p>25L</p> <p>500mL</p>
---------------	---

Directions for Use:	
---------------------	--

Restrains:	Re-treatment Interval: DO NOT retreat with BAYMEC LV Pour-On within the meat withholding period of 42 days.
------------	---

Contraindications:	
--------------------	--

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	<p>BAYMEC LV Pour-On is for external application to cattle only. The recommended dose rate for BAYMEC LV Pour-On is 1 mL/20 kg body weight. Apply along the topline of the animal. Cattle heavier than 650 kg should be dosed at 1 mL per 20 kg. A representative sample of animals should be weighed before treatment either with scales or a weigh band. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc.) DO NOT underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing. Rainfall on the herd within 2 hours of application may result in reduced efficacy of the product. Cattle should not be treated with BAYMEC LV Pour-On when the hide or hair is wet since the efficacy of the product may be reduced. Temporary hair loss occasionally occurs in the application area but will resolve without further treatment.</p>
----------------------------	--

General Directions:	NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.
---------------------	---

Withholding Periods:	<p>MEAT: DO NOT USE less than 42 days before slaughter for human consumption.</p> <p>MILK: ZERO (0) days.</p>
----------------------	---

Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product, confirm the current ESI from the manufacturer on 1800 995 709 or on the APVMA website (www.apvma.gov.au/residues).
---------------	---

Safety Directions:	Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves and goggles. Wash hands after use. After each day's use, wash gloves and goggles and contaminated clothing.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for a least 15 minutes and see a doctor. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.
First Aid Warnings:	
Additional User Safety:	
Environmental Statements:	<p>ENVIRONMENTAL PROTECTION</p> <p>Ivermectin is extremely toxic to aquatic species. DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used containers.</p>
Disposal:	<p>Containers up to 1 L: Dispose of container by wrapping with paper and putting in garbage.</p> <p>Containers over 1 L: Triple-rinse container into the medicated water, dip, drench, etc. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
Storage:	Store below 30°C (room temperature). Store in original container, tightly closed in a safe place.

Dose Table for Baymec LV Pour-On Broad Spectrum endectocide for Dairy and Beef Cattle 64610

Live weight kg	Dose Volume mL
51-100	5
101-150	7.5
151-200	10
201-250	12.5
251-300	15
301-350	17.5
351-400	20
401-450	22.5
451-500	25
501-550	27.5
551-600	30
601-650	32.5