



Product Name: BAYCOX COCCIDIOCIDAL SOLUTION  
APVMA Approval No: 45626/129927

Label Name:	BAYCOX COCCIDIOCIDAL SOLUTION
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS
Constituent Statements:	25 g/L TOLTRAZURIL
Claims:	For the treatment and control of coccidiosis in chickens caused by Eimeria species.  Baycox is added to the drinking water of chickens for the treatment and control of coccidiosis caused by Eimeria species including: Eimeria tenella, Eimeria necatrix, Eimeria acervulina and Eimeria maxima.
Net Contents:	100 mL 1 L 250 mL 5 L
Directions for Use:	
Restrictions:	DO NOT treat replacement pullets more than twice.
Contraindications:	
Precautions:	
Side Effects:	
Dosage and Administration:	Baycox is diluted at the rate of 3 litres per 1000 litres of drinking water (3:1000). It is administered on 2 consecutive days with an 8 hour treatment period on both days. Ensure

	<p>full lighting conditions during treatment. Do not withhold feed during treatment. Add the required amount of Baycox to the amount of water consumed in an 8 hour period and stir thoroughly. Allow for varying water consumption during hot weather. Treat the birds for 8 hours each day for 2 days (2x8 hour treatments). No other source of drinking water should be available to birds during the 8 hour treatment period.</p> <p>Broiler Treatment Programme: To use Baycox in a clinical outbreak obtain an accurate diagnosis. Begin treatment with Baycox as soon as coccidiosis is confirmed. A treatment period of 2 days is sufficient. If required, treatment may be repeated after 5 days.</p> <p>Broiler Preventive Programme: To use Baycox as a coccidiosis preventive in broiler production, give Baycox as above on days 9 and 10, 16 and 17, and 23 and 24 of life. Do not use a coccidiostat in the feed. An additional treatment may be necessary after 35 days of age but not within 14 days of slaughter.</p> <p>Layer Birds: Baycox may be used in replacement pullets and replacement layer breeder birds via the drinking water for an 8 hour period each day for two consecutive days. Treatment may be repeated after 5 days if required.</p>
--	--

General Directions:	<p>DO NOT USE GALVANISED OR RUSTY IRON WATER TANKS AND RETICULATION SYSTEMS.</p> <p>Baycox mixes with all types of water. However in areas where the drinking water is acidic, the water should be tested with pH test strips and brought to a minimum pH of 8.5 using sodium carbonate (washing soda).</p> <p>To avoid excessive foaming, add Baycox to the water after filling the supply tank.</p>
---------------------	---

Withholding Periods:	<p>MEAT: DO NOT USE less than 14 days before slaughter for human consumption.</p> <p>EGGS: DO NOT USE in laying hens. DO NOT USE in replacement pullets within 8 weeks of laying onset, where eggs are to be used or processed for human consumption. Eggs laid during the withholding period MUST NOT BE USED or processed for human consumption.</p>
----------------------	--

Trade Advice:	
---------------	--

Safety Directions:	Harmful if swallowed. Will irritate the eyes. Avoid contact with eyes.
--------------------	--

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
-------------------------	--

First Aid Warnings:	
---------------------	--

Additional User Safety:	Additional information is listed in the Safety Data Sheet.
-------------------------	--

Environmental Statements:	<p>Protection of Wildlife, Fish, Crustacea and Environment.</p> <p>Do not discharge waste water containing Baycox into dams, streams or watercourses.</p>
---------------------------	---

Disposal:	<p>[100 mL, 250 mL pack size]</p> <p>Dispose of empty container by wrapping with paper and putting in garbage.</p>
-----------	--

[1 L and 5 L pack sizes]

Triple-rinse container into the medicated water, dip, drench etc. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Storage:

Store below 30°C (room temperature) in the closed, original container, as cool as possible.