


Book Overlay 132 x 100 mm

Top

WARNING TOXIC, ECOTOXIC
Read label before use
Spinosad may cause target organ damage through prolonged or repeated exposure
Toxic to aquatic life with long lasting effects
Very toxic to terrestrial invertebrates



UN3082 Environmentally Hazardous substance, Liquid, N.O.S. (Spinosad)

Action: The active constituent in EXTINOSAD is a member of the "Spinozom" class of insecticides; a new generation insecticide with a unique mode of action.
Safety Precautions and Directions: Wash hands and exposed skin thoroughly after handling. Do not breathe mist or spray.
First Aid: Get medical advice if you feel unwell. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor.
Environmental Protection: Avoid release to the environment of the product or empty container.
Disposal: Preferably dispose of product by use. Otherwise dispose of product and packaging of an approved Agrecovery chemical disposal program. See www.agrecovery.co.nz for details. Triple rinse empty container. Add rinseate to dipwash. Recycle empty rinsed container at an approved Agrecovery recycling centre.
Emergency Response: In case of spillage wear appropriate protective clothing and prevent material from entering water ways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material.
Dispose of waste safely.
For specialist advice in an emergency call 0800 CHEMCALL (0800 243 622). If additional hazard information is required, refer to the Safety Data Sheet. For a copy please go to www.elanco.co.nz

PEEL HERE FOR LABEL DIRECTIONS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS
MEAT: NIL
MILK: Milk intended for human consumption must be discarded during treatment or for not less than 35 days following the last treatment

Extinosad: New science for high potency, clean wool and operator safety.

- NO WOOL WITHHOLDING PERIOD
- NO MEAT WITHHOLDING PERIOD
- EXTINOSAD is exempt from the Wool Pro dip withholding times for all fleeces types.



Storage: Store below 30 °C in a dry place in a tightly closed container. Do not freeze. Do not store in direct sunlight.

Registered to:
Elanco Australasia Pty Ltd
27/88 Shortland Street, Auckland, 1010, New Zealand
HELPLINE: Phone 0800 426 633
www.elanco.co.nz

Registered pursuant to the ACVM Act 1997, No A000206
See www.foodsafetynz.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996, No HSR001884
See www.sgs.govt.nz for approval conditions

BATCH NUMBER

EXPIRY

YL402409A
AH0492

135 mm

130 mm

Base


MIXING DIRECTIONS: LICE AND FLY
SHAKE CONTAINER SEVERAL TIMES BEFORE USE
This product should be mixed at the rate specific for the desired use as shown in the Directions for Use tables. Mix well, preferably by re-circulating the dip through the pump before use. If using shower dips or hand jetting wands. If using plunge dips, ensure the even distribution and mixing of Extinosad throughout the dip. For best results, make an initial dilution by adding the concentrate to a bucket of water. Mix well and pour along the length of a near empty dip sump. Fill with water and mix the dip sump contents well before use. Prepare fresh dip, jetting fluid or fly-strike dressing daily. At the end of the day, discard unused product. It is good agricultural practice not to treat animals when they are wet or if rain is likely. However, even moderate to heavy rainfall following treatment does not adversely affect the efficacy of the product.



For advice on your dipping programme, contact your supplier or call the Elanco HELPLINE: 0800 426 633

YL402409A

132 mm

107 mm

PRODUCT INFO		
BLUE #: 402409A	Product Name: Extinosad	
Item Code: YL402409A	Component: Booklet Label	
Product Code: AH0492	Pack Size: 10L	
Previous Item Code: YL402107X		
ARTWORK INFO		
Template: YL_130x135mm_v2	Packaging Specs: N/A	
BarcodeType: EAN 13: 9418933049231	Add. Info: N/A	
	Minimum Core Data Point Size: 6pt	
GTIN: GTIN Not required	Proof #	PS) By/Date MFA 16-AUG-2019
		
ELANCO ARTWORK LEGEND v13		

KEY	
	Text Area
	Variable Data and Code Area

© Elanco Australasia Pty Ltd

132 mm

100 mm

1

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

DIRECTIONS FOR USE: LICE
For the treatment of lice on long wool Merino sheep (greater than 3 months wool), including strains resistant to synthetic pyrethroids.
For the control of lice on coarse wool breeds, including strains resistant to synthetic pyrethroids.

FOR THE CONTROL OF LICE ON COARSE WOOL BREEDS

TREATMENT METHOD	DOSE	COMMENTS
PLUNGE	Initial charge: Use 1: 2500 dilution e.g.: 400 mL dip in 1000 L water. Replenishment rate: Use 1: 1700 dilution e.g.: 600 mL dip in 1000 L water.	Plunge dip each animal for at least 20 seconds ensuring the head is immersed twice. Ensure swim length is > 7.0 metres. For best results use constant replenishment.
CONTINUOUS REPLENISHMENT (CR) SHOWER	Use rate: Use 1: 1250 dilution e.g.: 800 mL dip in 1000 L water.	Shower: For best results using shower dip equipment, allow sufficient time in the dip to achieve thorough wetting. Do not allow the sump to fall below ¾ of total volume and for best results use constant replenishment.
STANDARD SHOWER	Initial charge: Use 1: 1250 dilution (20 ppm) e.g.: 800 mL dip in 1000 L water. Replenishment rate: Use 1: 625 dilution (40 ppm) e.g.: 1.6 L dip in 1000 L water.	Do not allow the sump to fall below ¾ of total volume and for best results use constant replenishment.

2

FOR THE TREATMENT OF LICE ON LONG WOOL MERINO SHEEP

TREATMENT METHOD	DOSE	COMMENTS
LICE JETTING	Use 1: 1000 dilution e.g.: 1 L dip in 1000 L water.	Jet: Using jetting equipment, apply a minimum of 2 litres per sheep or 500 mL of fluid for each month of wool growth, up to a maximum of 5 litres per sheep. Apply from the poll to the tail base, around the neck and along each side saturating to skin level. Lousy long wool Merino sheep treated 6 months or more before shearing may require a second treatment.
SHOWER	Initial charge: Use 1: 1000 dilution e.g.: 1 L dip in 1000 L water. Replenishment rate: Use 1: 500 dilution e.g.: 1 L dip in 500 L water.	Shower: For best results using shower dip equipment, allow sufficient time in the dip to achieve thorough wetting. Do not allow the sump to fall below ¾ of total volume and for best results use constant replenishment. Lousy long wool Merino sheep treated 6 months or more before shearing may require a second treatment.

GENERAL COMMENTS
For effective control of lice, thorough wetting is essential, check skin wetting using a wetting pencil. For best results, clean and refill the sump after one sheep per 2 litres of initial sump volume have been dipped.
Optimum lice control requires the application of good agricultural practice. This includes attention to stock security following treatment, quarantine treatment of new stock arrivals, thorough and accurate dipping management and utilising the principles of Integrated Pest Management recommended by Wool Pro.

3

DIRECTIONS FOR USE: FLY
For the treatment of and short-term prevention of blowfly strike (*Lucilia cuprina*, *Lucilia sericata* and secondary strike sp.) on sheep, including strains resistant to organophosphates.
FOR THE TREATMENT OF AND SHORT-TERM PREVENTION OF BLOWFLY STRIKE ON ALL BREEDS OF SHEEP

TREATMENT	RATE	COMMENTS
FLY JETTING	Use 1: 1000 dilution e.g.: 1 L dip in 1000 L water.	When applied as directed, this product will protect sheep against fly-strike for 2-4 weeks. Using a hand-gun designed specifically for jetting, apply a minimum of 2 litres per sheep or 500 mL of fluid for each month of wool growth, up to a maximum of 5 litres per sheep. Apply from the poll to the tail base and around the breech and pizzle as appropriate. Use Pressure requirements: 500-700 kPa (75-100 psi).
FLY STRIKE DRESSING	Use 1: 1000 dilution e.g.: 5 mL dip in 5 L water.	Prepare a fresh dilution of fly-strike dressing each day. Remove the wool from around the wound using clippers or shears. Apply 1 to 2 litres Extinosad onto the wound and ensure the maggots are destroyed.

4

Booklet pages must be in multiples of 4.
You may delete or add pages in multiples of 4 to fit artwork.