

PRODUCT PROFILE

Monensin with Microtracer®

RUMENSIN®

PREMIX FOR USE IN BEEF CATTLE



Please note there are additional indications for use in dairy cattle; please refer to the full label.

INDICATIONS:

1. For improved feed efficiency in beef cattle (steers and heifers) fed in confinement for slaughter.
2. As an aid in the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in cattle. Note to user: Coccidiosis occurs sporadically in first lactation dairy heifers, but is not considered a significant disease in mature dairy cows.
3. For increased rate of weight gain in growing cattle on pasture (slaughter, stocker and feeder cattle, and beef and dairy replacement heifers) of greater than 180 kg (400 lb) body weight.

FEEDING DIRECTIONS

Complete Diet: which includes complete feed plus roughage



IMPORTANT: Must be thoroughly mixed in feeds before use. Feed continuously at a rate of not less than 33 g and not more than 48 g monensin activity per tonne (1,000 kg) until animals reach market weight.



Feed continuously at a rate of 22 g monensin activity per tonne (1,000 kg) during periods of exposure to coccidiosis or when coccidiosis is likely to be a hazard.



Hand feed continuously a minimum of 0.5 kg of medicated supplement per head per day, to supply 200 mg of monensin per head per day. The medicated supplement must be hand fed from the beginning to the end of the pasture season.

WARNINGS:

1. For three indications listed above. Do not supplement monensin from other sources (eg. other feedstuffs containing monensin, the Rumensin Controlled Release Capsule or Kexxtone).
2. No withdrawal period is required when used according to the label.
3. When mixing and handling Rumensin Premix, avoid inhalation, oral exposure and direct contact with skin or eyes. Use protective clothing, impervious gloves and dust mask. Operators should wash thoroughly with soap and water after handling.
4. Keep out of reach of children.

CAUTIONS:

1. Do not exceed recommended levels as reduced average daily gains may result.
2. Do not allow canines, horses, other equines or guinea fowl access to formulations containing monensin. Ingestion of monensin by these species has been fatal.
3. Do not use Rumensin medicated feed for the treatment of outbreaks of coccidiosis.
4. May be used in feeds containing the pellet-binding agents, Bentonite (2%), Attapulgit (2%), Kaolin (2.5%), Lignin Sulfonate (4%), Carboxymethylcellulose (0.1%) or Agri-Colloid.
5. NOT TO BE USED AFTER EXPIRATION DATE. Do not use thixotropic supplement after eight weeks storage.

The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.