## Parvovirus Vaccine Killed Virus Erysipelothrix RhusiopathiaeLeptospira CanicolaGrippotyphosa-HardjoIcterohaemorrhagiaePomona Bacterin

For Use In Animals Only

Parvo Shield™ L5E

**INDICATIONS:** For use in healthy swine as an aid in the prevention of disease caused by porcine parvovirus, *Erysipelothrix rhusiopathiae*, *Leptospira canicola*, *grippotyphosa*, *hardjo*, *icterohaemorrhagiae*, and *pomona*. This product has not been tested in pregnant animals.

**ADJUVANT:** Aluminum hydroxide.

**DIRECTIONS:** Shake well before using. Administer 5 mL intramuscularly to gilts and sows 4-6 weeks prior to breeding. Repeat in 3-4 weeks. Revaccinate with a single dose 4-6 weeks prior to each subsequent breeding. Vaccinate boars semiannually.

**PRECAUTIONS:** Store out of direct sunlight at 2°-8°C (35°-46°F). DO NOT FREEZE.

Do not mix with other products except as specified on this label. In case of human exposure contact a physician. Use entire contents when first opened.

Do not vaccinate within 21 days prior to slaughter. Anaphylactic reactions may occur.

Symptomatic treatment: Epinephrine. Contains amphotericin B, gentamicin, and thimerosal as preservatives.

VLN/PCN: 196/48C5.20

Elanco US Inc., Fort Dodge, IA 50501 USA; 1-800-428-4441

Parvo Shield, Elanco and the diagonal bar logo are trademarks

of Elanco or its affiliates.



YL101164A W1c