



Product Name: Elanco Viper Pour-On Lousicide for Sheep  
APVMA Approval No: 87489/130385

Label Name:	Elanco Viper Pour-On Lousicide for Sheep
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS
Constituent Statements:	10 g/L THIAACLOPRID
Claims:	For the control of neonicotinoid susceptible and IGR resistant strains of body lice ( <i>Bovicola ovis</i> ) when applied to shorn sheep up to 7 days off shears and unshorn lambs from 2 weeks to 2 months of age. Protects sheep from re-infestation with lice for 4 weeks after treatment when applied within 24 hours of shearing.  Resistance may develop to any chemical.  READ DIRECTIONS FOR USE BEFORE OPENING OR USING THIS PRODUCT
Net Contents:	120 L 110 L 100 L 60 L 55 L 22 L 20 L 5 L
Directions for Use:	
Restrains:	DO NOT USE more than 7 days after shearing DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.
Contraindications:	

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	This section contains file attachment.
----------------------------	--

General Directions:	<p>Apply product only with the Elanco 6-stripe applicator.</p> <p>Sheep must be treated within 7 days of shearing. Efficacy is not affected by moderate rainfall either before or after treatment (50 mm rain 1 hour prior to treatment and/or 12.5 mm 2 hours after treatment and/or 25 mm 24 hours after treatment). Treated sheep must not be mixed with lousy sheep. This product should not be used on sheep which have not been cleanly shorn for any reason e.g. mycotic dermatitis.</p> <p>Management practices which promote good lice control such as complete musters, accurate treatment of every animal according to label, ensuring all animals can be treated at the same time, no split shearings, stock-proof fencing to prevent external introductions, quarantine and lice treatment of all newly purchased stock are essential for effective lice control.</p> <p>This product may be used in conjunction with worm control drenches and at the same time as vaccination.</p> <p>This product contains a blue dye to identify treated sheep. This dye is scourable from the wool.</p> <p>This product is safe for use on breeding ewes and rams.</p>
---------------------	--

Withholding Periods:	<p>MEAT: DO NOT USE less than 42 days before slaughter for human consumption.</p> <p>MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.</p> <p>WOOL: DO NOT USE less than 6 months before shearing or fibre collection</p>
----------------------	---

Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 56 days before slaughter for export. Before using this product, confirm the current ESI from Elanco Australasia Pty Ltd on 1800 995 709 or APVMA website (<a href="http://www.apvma.gov.au/residues">www.apvma.gov.au/residues</a>).</p>
---------------	---

Safety Directions:	<p>Will damage eyes. Avoid contact with eyes. When using the product, wear cotton overalls over normal clothing, buttoned to the neck and wrist and a washable hat, PVC or rubber apron and chemical-resistant gloves and water resistant footwear and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.</p>
--------------------	--

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126</p>
-------------------------	--

First Aid Warnings:	
---------------------	--

Additional User Safety:	<p>Sheep Rehandling Interval: Four weeks. If sheep must be handled during this interval, cotton overalls buttoned to the neck and wrist (or equivalent clothing) should be worn.</p> <p>Additional information is listed in the Safety Data Sheet.</p>
-------------------------	--

Environmental Statements:	<p><b>ENVIRONMENTAL PROTECTION:</b> Very toxic to aquatic life. DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used container.</p>
---------------------------	--

Disposal:	<p>Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.</p>
-----------	---

Storage:	<p>Store below 30°C (Room temperature). Store in original container, tightly closed in a safe place. Protect from light.</p>
----------	--

## DOSAGE AND ADMINISTRATION:

### SHAKE WELL BEFORE USE

#### Critical comments:

Use only on sheep that have been cleanly shorn, or unshorn lambs from 2 weeks to 2 months of age.

Avoid treating ewes less than 6 weeks before lambing commences or while ewes are lambing or have lambs less than 2 weeks old at foot, as live lice can persist and infest lambs.



Treated sheep should not be mixed with untreated sheep until 6 weeks after treatment.

Apply the entire dose to sheep under 30kg as a single broadband stripe from poll to tail butt.

For sheep over 30kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the butt of the tail. The stripes should be no further than 10 cm apart.

Refer to the diagrams below.

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within the group (e.g. lambs), dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Bodyweight (kg)	Dose (mL)	Method	Apply in the pattern below
6.1 – 8.0	12	Single stripe	
8.1 – 10.0	15		
10.1 – 12.5	20		
12.6 – 15	25		
15.1 – 20	30		
20.1 – 30	40		
30.1 – 55	60	Double stripe	
55.1 – 80	80		
80.1 – 100	90		
Add 10 mL for each 10 kg above 100 kg			

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**