

FOR EXTERNAL ANIMAL USE ONLY / SLEGS VIR UITWENDIGE DIERGEBRUIK

DRASTIC Deadline®



Reg. No./ -nr G723 (Act/Wet 36/1947)
Namibia/Namibië: NSR 5

Ectoparasiticide for instant use on cattle, sheep, ostriches and game. Ektoparasietmiddel gereed vir gebruik op beeste, skape, volstruise en wild.


- Controls ticks (including Karoo Paralysis ticks) on cattle, sheep, ostriches and game
- Controls tsetse fly worry on cattle
- Controls feather mites on ostriches
- Controls red lice on cattle
- Sterilises female ticks
- Easy application onto skin
- Residual action
- Non-systemic action
- Has a 21 day residual action against ticks on ostriches
- Beheer bosluise (insluitend Karooverlammingsbosluise) op beeste, skape, volstruise en wild
- Beheer tsetsevlieg plaie by beeste
- Beheer veermyte op volstruise
- Beheer rooiluise op beeste
- Steriliseer wyfiebosluise
- Maklike aanwending op die vel
- Nawerkende effek
- 'n Nie-sistemiese werking
- Het 'n 21 dae nawerkende effek op bosluise op volstruise

CAUTION / VERSIGTIG

Store in a cool, dry place / Berg in 'n koel, droë plek

Contains / Bevat:
Flumethrin / Flumetrien 1 % m/v

1e



For cattle, sheep, ostriches and game
Vir beeste, skape, volstruise en wild

WARNINGS: Before using this product read the label carefully. No withdrawal period for milk or meat. Keep out of reach of children, uninformed persons and animals. Store in a safe place away from feed and food. Toxic to bees and fish. Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder. Skin reaction may occur in a small percentage of cases, if this happens wash the affected animals with soap and water.

DIRECTIONS FOR USE: Use only as directed. Shake well before use. Treat at regular intervals, according to tick challenge.
The basic dosage is 1 ml per 10 kg body mass.

For tsetse flies treat every 14 days according to fly challenge.

Cattle: Apply the required dose volume along the backline.

Sheep: Apply the required dose volume in 4 parts to be the bare skin areas of the groin and axillae. In the case of sheep infested with Karoo Paralysis Tick (*Ixodes rubicundus*), Drastic Deadline has to be applied early in the season, when these ticks are first observed (either on the grazing, or on the animals). Two to three treatments at 7 to 10 day intervals at this time should suffice to protect sheep against Karoo Paralysis, depending on the severity of the tick challenge.

Ostrich: Apply the required dose volume along the backline.

Game: Apply the required dose volume with a suitable applicator.

If a self-medicating device is used, it is important that the device only remains activated for as long as it takes to deliver one dosage to the animals in the particular paddock/camp, and that the device is not used for continuous low-level treatment.

DOSAGE SCHEME Up to	DOSE
100 kg body mass	10 ml
200 kg body mass	20 ml
300 kg body mass	30 ml
400 kg body mass	40 ml
500 kg body mass	50 ml

Squeeze bottle with dosemeter. Squeeze required volume into dosemeter.

LIVE MASS IN KG	NUMBER OF DOSES PER 1 LITRE
100 kg body mass	100
200 kg body mass	50
300 kg body mass	33
400 kg body mass	25
500 kg body mass	20

Registration Holder/Registrasiehouer:
Bayer (Pty) Ltd., Animal Health Division,
Bayer (Edms) Bpk., Diergesondheidsafdeling,
Reg. No./ -nr.: 1968/011/92/07
27 Wrench Rd/Wrenchweg 27, Isando, 1601, R.S.A.
Tel: (011) 921 5736, Fax/Faks: (011) 921 5751
www.bayeranimalhealth.co.za

© is the registered trademark of Bayer AG, Germany
© is die geregistreerde handelsmerk van Bayer AG, Duitsland

WAARSKUWINGS: Lees die etiket deeglik voordat die produk gebruik word. Geen onttrekkingsperiode vir melk en vleis nie. Hou buite bereik van kinders, oningeligte persone en diere. Berg op 'n veilige plek we van voedsel en voer. Giftig vir bye en visse. Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer. Vel reaksies mag in 'n klein persentasie gevalle voorkom, indien dit sou gebeur, was die geaffekteerde diere met seep en water.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui. Skud goed voor gebruik. Behandel met gereelde tussenposes, na geland van bosluislading.

Die basiese dosis is 1 ml per 10 kg liggaamsmassa.

Vir tsetsevlieë behandel met 14 dae tussenposse,

na geland van vliegplaag.

Beeste: Wend die verlangde dosis al met die ruglyn langs aan.

Skape: Wend die verlangde dosis in 4 dele aan op die kaal huid van die lieste en oksels (onder die blaaië). In die geval skape wat blootgestel is aan besmetting met Karooverlammingsbosluise (*Ixodes rubicundus*), moet Drastic Deadline vroeg in die seisoen, wanneer die eerste bosluise gewaar word (hetsy op die weiding, of op die skape), aangewend word.

Twee tot drie behandelings met 7 tot 10 dae tussenposes op hierdie tyd behoort voldoende beskerming teen Karooverlamming te bied, afhangend van die graad van die bosluisbesmetting.

Volstruise: Wend die verlangde dosis aan al met ruglyn langs.

Wild: Wend die verlangde dosis met 'n geskikte aanwender aan. Indien 'n selftoediener gebruik word, is dit belangrik dat die toestel geaktiveer bly slegs vir solank as was nodig is om die diere in die betrokke kamp eenmaal te behandel en dat die toestel nie gebruik word vir langdurige laevlakbehandeling nie.

DOSISTABEL tot	DOSIS
100 kg liggaamsmassa	10 ml
200 kg liggaamsmassa	20 ml
300 kg liggaamsmassa	30 ml
400 kg liggaamsmassa	40 ml
500 kg liggaamsmassa	50 ml

Druk bottel met maatbeker. Druk benodigde hoeveelheid in maatbeker in.

LEWENDE MASSA IN KG	AANTAL DOSISSE PER 1 LITRE
100 kg liggaamsmassa	100
200 kg liggaamsmassa	50
300 kg liggaamsmassa	33
400 kg liggaamsmassa	25
500 kg liggaamsmassa	20

Lot nr./ No:
Verw./Manuf:
Exp:

Bayer

58183672 MD