

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



Monepantel is a member of the Amino-Acetonitrile Derivatives (AADs) class of anthelmintics and can be used for the treatment and control of AAD-sensitive strains of gastro-intestinal roundworms (nematodes), including macrocyclic lactone, benzimidazole (white), levamisole and morantel (clear) and closantel-resistant strains in sheep.

ELANCO AUSTRALASIA PTY LTD
7 Eden Park Drive, Macquarie Park, NSW, 2113
TEL: Toll free 1800 226 324

www.elanco.com.au



**Broad Spectrum Oral Anthelmintic
for Sheep**

**ACTIVE CONSTITUENT:
25 mg/mL Monepantel**



10 L



x 4000

YL402887A 311250

READ THE ATTACHED LEAFLET FOR FULL INSTRUCTIONS

DIRECTIONS FOR USE

Restraints: DO NOT USE in sheep less than 10 kg body weight. DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

Re-treatment Interval: DO NOT re-treat animals for 42 days after the last treatment.

Contraindications: This product is contraindicated for use in lambs under 6 weeks of age or less than 10 kg body weight.

Dosage and administration: Use remaining product within 12 months of opening the container.

ZOLVIX PLUS is a ready-to-use oral solution.

Check dose rates and drench gun before treatment commences.

The dose is 0.5 mL ZOLVIX PLUS /5 kg live weight (1 mL/10 kg), equivalent to 2.5 mg monepantel/kg and 0.2 mg abamectin/kg live weight. Drench sheep orally using an Elanco recommended drench gun. Maintain applicator gun carefully to ensure accurate dosage. After use, clean gun by flushing with warm, soapy water. Rinse with cold water.

Body weight (kg)	Dose (mL)	No. Treated/ 10 L
10–15	1.5	6666
16–20	2.0	5000
21–25	2.5	4000
26–30	3.0	3332
31–35	3.5	2856
36–40	4.0	2500
41–50	5.0	2000
51–60	6.0	1666
61–70	7.0	1428
71–80	8.0	1250

Sheep in excess of 80 kg body weight should be dosed at 1 mL/10 kg. A representative sample of animals should be weighed before treatment.

Dose the mob to the heaviest sheep by live weight in each group (ewes, wethers, rams, lambs). DO NOT under dose. Where there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing. A correct drenching technique is recommended; carefully place the drench gun nozzle in the side of the mouth and over the back of the tongue before delivering the dose.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 14 days before slaughter for human consumption.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. Before using this product, confirm the current ESI from Elanco Australasia on 1800 226 324 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS: Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. When opening the container and using the product, wear elbow length chemical-resistant gloves. If product on skin, immediately, wash area with soap and water. Wash hands after use. After each day's use wash gloves.

FIRST AID: If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

ENVIRONMENTAL PROTECTION: Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the *drumMUSTER* logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not

recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE: Store below 30°C (room temperature). Store in the original tightly closed container. Protect from light.

Customer Information Australia: 1800 226 324

APVMA Approval No. 69763/134202

SAFETY DATA SHEET: If additional hazard information is required, refer to the Safety Data Sheet. For a copy please go to www.elanco.com.au



Batch No.	
Expiry Date	

READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

Zolvix contains monepantel, a member of the Amino-Acetonitrile Derivatives (AADs) class of anthelmintics. When used at the recommended dose rate (2.5 mg/kg), it is effective against AAD-sensitive strains of the following gastro-intestinal roundworms (nematodes) including those resistant to macrocyclic lactones, beimidazoles (white drenches), levamisole and morantel (clear), and closantel (*Haemonchus contortus* only) in sheep. Resistance may develop to any chemical.

Efficacy spectrum

Barber's pole worm *Haemonchus contortus***,
Small brown stomach worm *Ostertagia (Teladorsagia) circumcincta**,
*Ostertagia (Teladorsagia) trifurcata**, *Ostertagia (Teladorsagia) davtiani**,
Ostertagia (Teladorsagia) spp. (inhibited L4),
Black scour worm *Trichostrongylus colubriformis**,
*Trichostrongylus vitrinus**, *Trichostrongylus rugatus**,
Stomach hair worm *Trichostrongylus axei**,
Thin necked intestinal worm *Nematodirus filicollis**,
Nematodirus spathiger (adult only),
Small intestinal worm *Cooperia curticei**, *Cooperia oncophora**,
Large mouthing bowel worm *Chabertia ovina**,
Large bowel worm *Oesophagostomum venulosum* (immature L4 only).

* Includes **adult and immature L4 stage**

** Includes **adult, immature L4 stage and inhibited L4 stage**

Ask your veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted before any parasite treatment is used

READ THE ATTACHED LEAFLET FOR FULL INSTRUCTIONS

DIRECTIONS FOR USE

Restraints:

DO NOT USE in sheep less than 10 kg body weight.

DO NOT USE in female sheep which are producing or may in the future produce milk that may be used or processed for human consumption.

Re-treatment interval: DO NOT re-treat less than 21 days after the last treatment.

Dosage and administration

Use remaining product within 12 months of opening the container.

ZOLVIX is a ready-to-use solution. Check dose rates and drench gun before treatment commences. The dose is 0.5 mL ZOLVIX/5 kg live weight (1 mL/10 kg), equivalent to 2.5 mg MONEPANTEL/kg live weight. Drench sheep orally, using an Elanco recommended drench gun. Maintain applicator gun carefully to ensure accurate dosage. After use, clean gun by flushing with warm, soapy water. Rinse with cold water.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 14 days before slaughter for human consumption.

MILK: DO NOT USE in female sheep which are producing or may in the future produce milk that may be used or processed for human consumption.

TRADE ADVICE-Export Slaughter Interval

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 14 days before slaughter for export. Before using this product, confirm the current ESI from Elanco Australasia Pty Ltd on 1800 226 324 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS

May irritate the eyes. Will irritate the skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. When using the product wear elbow length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 1126).

ENVIRONMENTAL PROTECTION

Do not contaminate dams, rivers, streams or other waterways with the chemicals or used containers.

DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30 °C (room temperature). Store in the original tightly closed container. Use remaining product within 12 months of opening the container.

Elanco Australasia Pty Ltd
7 Eden Park Drive,
Macquarie Park, NSW, 2113
TEL: Toll free 1800 226 324
Manufactured in New Zealand



5 420036 907664

Batch No.	
Expiry Date	