

20 Gilt/Sow Doses
50/100 Piglet Doses
100 mL

Bordetella
Bronchiseptica-Erysipelothrix
Rhusiopathiae-Pasteurella
Multocida Bacterin-Toxoid

Elanco



For Use In Animals Only

INDICATIONS: This product has been shown to be effective for vaccination of healthy swine, 7 days of age or older, against atrophic rhinitis caused by *Bordetella bronchiseptica* or the toxin of *Pasteurella multocida* Types A and D, erysipelas caused by *Erysipelothrix rhusiopathiae*, and pneumonia caused by *Pasteurella multocida* Type A. The duration of immunity for this product has not been established. For more information regarding efficacy and safety data, see productdata.aphis.usda.

ADJUVANT: Aluminum hydroxide.

DIRECTIONS: Shake well before using. Administer intramuscularly or subcutaneously. Four doses must be given (two gilt/sow doses during pregnancy and two piglet doses). Vaccinate gilts and sows with a 5 mL dose at 5 and 2 weeks prior to farrowing. Vaccinate piglets from vaccinated dams with a 1 mL dose at 7 to 10 days of age and a 2 mL dose 2 weeks later. The need for annual booster vaccinations has not been established for this product; consultation with a veterinarian is recommended. For advice on revaccination frequency, in general or in the face of maternal antibody, stress, or a disease exposure, consultation with a veterinarian is recommended.

PRECAUTIONS: Store out of direct sunlight at 2 to 8°C (35° to 46°F). DO NOT FREEZE. Do not mix with other products, except as specified on this label. Use entire contents when first opened. Do not vaccinate within 21 days prior to slaughter. In case of human exposure, contact a physician. Anaphylactic reactions may occur. Symptomatic treatment: Epinephrine.

VLN/PCN 196/7054.01

Elanco US Inc., Fort Dodge, IA 50501 USA; 1-800-428-4441

Rhini Shield, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.