


WARNING
KEEP OUT OF REACH OF CHILDREN
READ LABEL BEFORE USE
FOR ANIMAL TREATMENT ONLY

Elanco™

Bayticol™**Pour-on Tickicide**

Active Constituent:
10 g/L flumethrin

1 LITRE

Handling Precautions: Bayticol may cause a temporary skin irritation in some individuals if applied to the skin. Presumed to/may cause organ damage (dermatotoxic) from repeated exposure at high doses. Avoid contact with eyes, mouth, hands and skin. Wear protective gloves (e.g. disposable nitrile safety gloves) and protective clothing when handling. Additional protective clothes (e.g. long sleeved shirt, long pants, boots and water resistant apron) are necessary if an amount of more than 10 litres of product is applied to animals per day. Do not drink, eat or smoke while using the product. Wash hands, exposed skin and face before breaks and at the end of work. Do not breathe vapour. Avoid ingestion.

First Aid: If on skin remove contaminated clothing and wash skin thoroughly with soap and water. In case of accidental spillage into the eyes or mouth wash with plenty of water. If on skin wash off with soap and water. If you feel unwell get medical advice. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

Environmental Protection: Harmful to fish and aquatic life with long lasting effects. Avoid contamination of any water supply with product or empty container. Harmful to terrestrial invertebrates. Avoid release to the environment.

Emergency Response: For specialist advice in an emergency call 0800 734 607(24 hours)

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

- Apply Bayticol Pour-On evenly along the midline of the back from the front of the shoulder to the tail head.
- Use a plastic measuring cup or automatic drenching gun to apply.
- Control of ticks is not affected by rain falling within a few hours of treatment.

CATTLE	DOSE VOLUMES
Up to 200 kg liveweight	20 mL
200–300 kg	30 mL
301–400 kg	40 mL
Over 400 kg	10 mL/100kg

- Bayticol spreads through the oils of the skin and hair coat to all parts of the animal's body. Behaviour such as grooming, tail swishing and scratching, aid the movement of active ingredient to the extremities of the body which are often affected by ticks.
- Bayticol is active against all 3 stages of the New Zealand cattle tick (*Haemaphysalis longicornis*) lifecycle. It also causes infertility of ticks which survive treatment, by inhibiting egg laying or rendering eggs sterile. This action helps reduce contamination of pastures.

DEER	DOSE VOLUMES
Up to 6 months	10 mL
Yearlings and adults	20 mL

- Retreatment should be undertaken when ticks start to reappear. Overseas trials with various tick species indicate that Bayticol provides protection against reinfestation for 3 to 6 weeks.
- For optimum control in deer, breeding hinds should be treated every three weeks from August to fawning. This programme will reduce the number of nymphs reaching the adult stage, which infest fawn during November–December. Treatment of hinds can be continued after fawning if this is practical. Other stock should be treated every three weeks from August, or as required.

ADVERSE EFFECTS

- Signs of mild irritation may be observed for 24–48 hours after treatment.
- Hair loss at the site of application may occur on deer which are shedding their winter coat at the time of treatment. This effect is temporary and can be minimised by spreading the dose over the entire backline. Growth of summer coat is not affected.

WITHHOLDING PERIOD

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

There is NO WITHHOLDING PERIOD for meat and milk from treated animals.

STORAGE

Once opened, product can be used for up to 9 months. Store locked up below 30°C (room temperature). Do not freeze. Store away from food and feed.

DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill/ facility. Triple rinse container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at an approved landfill/facility. Do not use container for any other purpose.

Registered pursuant to the ACVM Act 1997, No. A005335
See www.foodsafety.govt.nz for registration conditions.

Elanco New Zealand
106 Wiri Station Road, Manukau, Auckland 2104.
Customer Information Line 0800 446 121.

Bayticol is sold by Elanco or its affiliates and is not a product of Bayer. The Product Name Bayticol is owned by Bayer and used under license.


© 2022 Elanco or its affiliates



Batch:
Expiry:

5 420036 980193

Note: Pharmacode can be moved left or right along bottom edge as long as print free area either side is maintained.

**PRODUCT INFO**

BLUE #:	502243APK	Product Name:	Bayticol 1% Pour On
Item Code:	90209975	Component:	Label
Product Code:	LBAH	Pack Size:	1L
Previous Item Code	90207316		

ARTWORK INFO

Template:	LA_280x120mm_v3b	Packaging Spec(s):	TPM_016173
Barcodes/Type:	EAN 13: 5420036980193 Laetus : 43	Add. Info:	EAN13 code added to label
GTIN:	N/A	Proof #	P1a
		By/Date	RA 22-JUN-2022
NON PRINTING	PMS 185 C	BLACK	MAGENTA
		YELLOW	CYAN
			PMS 285 C

ELANCO ARTWORK LEGEND v16

LEGEND KEY

Template: LA_280x120mm_v3b

Text Free Area	Print Free Area
Variable Data and Code area	
50 mm Rule Bar	

- Item code placement is flexible and can be moved to suit artwork
- GTIN code placement is flexible but where possible should be placed close to the variable data info
- Ensure VD Headers remain RIGHT aligned and that the longest Header Text starts 2mm from the edge
- For 3 VD Headers - Use Lines (2, 3 & 4)
- For 2 VD Headers - Use Lines (3 & 4)
- For 1 VD Header - Use Line (4)
- Adjust VD Print Free area to the required number of headers as indicated by the dotted lines.

20mm min.
Note: A minimum of 20mm is required for Print Free VD area. This can be extended as required. VD Headers can be any length.