

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS  
FOR ANIMAL TREATMENT ONLY



# VIPER<sup>®</sup>

Pour-On

Lousicide for Sheep

ACTIVE CONSTITUENT: 10g/L THIACTOPRID

For the control of neonicotinoid susceptible and IGR resistant strains of body lice (*Bovicola ovis*) on shorn sheep up to 7 days off shears. This product can be expected to protect sheep against re-infestation with lice for 4 weeks after treatment when applied within 24 hours of shearing.



20L



18759 06/16

WhiteBlack

Operator:		Proof Name:	
Product Code:		Folio type:	
Master Card:			
Varnish Type 1:			
Varnish Type 3:			



**POUR-ON LOUSICIDE FOR SHEEP**

**READ DIRECTIONS FOR USE BEFORE OPENING OR USING THIS PRODUCT.**

VIPER® Pour-On is for the control of neotectoid susceptible and IGR resistant strains of body lice (*Bovicola ovis*) on shorn sheep up to 7 days off shears.

**DIRECTIONS FOR USE**

**RESTRAINTS: DO NOT USE** more than 7 days after shearing. **DO NOT USE** on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

**SHAKE WELL BEFORE USE**

**DOSAGE AND ADMINISTRATION**  
**Critical comments:** Use only on sheep that have been cleanly shorn. Avoid treating ewes less than 6 weeks before lambing commences, while ewes are lambing or have lambs at foot, as live lice can persist and infect lambs. Treated sheep should not be mixed with untreated sheep until 6 weeks after treatment. Apply the entire dose to sheep under 30kg as a single broadband stripe from poll to tail butt. For sheep over 30kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the butt of the tail. The stripes should be no further than 10cm apart. Refer to the diagrams below.

Bodyweight (kg)	Dose (mL)	Method	Apply in the pattern below
5.0 – 6.0	8		
6.1 – 8.0	12		
8.1 – 10.0	15	Single stripe	
10.1 – 12.5	20		
12.6 – 15	25		
15.1 – 20	30		
20.1 – 30	40		
30.1 – 55	60		
55.1 – 80	80	Double stripe	
80.1 – 100	90		
Add 10mL for each 10kg above 100kg.			

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHOLDING PERIODS**

**MEAT: DO NOT USE less than 42 days before slaughter for human consumption.**  
**MILK: DO NOT USE on ewes which are producing, or may in the future produce, milk or milk products for human consumption.**  
**WOOL: DO NOT USE less than 6 months before shearing or fibre collection.**

**TRADE ADVICE**  
**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 56 days before slaughter for export.**

**PEEL HERE**

**SAFETY DIRECTIONS**  
 Will damage eyes. Avoid contact with eyes. When using the product, wear cotton overalls over normal clothing, buttoned to the neck and wrist and a washable hat, PVC or rubber apron and rubber gloves. If gloves are used, wash them thoroughly with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**Sheep Rehoming Interval: Four weeks. If sheep must be handled during this interval, cotton overalls buttoned to the neck and wrist (or equivalent clothing) should be worn.**

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Additional information is listed in the Safety Data Sheet.

**THE ENVIRONMENT**

Very toxic to aquatic life. DO NOT contaminate streams, rivers, wetlands or watercourses with this product or used containers.

**DISPOSAL**

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. DO NOT dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured container in a safe place, away from the surface in a deep pit, preferably at least 50cm deep, for this purpose. Clear of weeds, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

**STORAGE**  
 Store below 30°C (room temperature). Store in original container, tightly closed in a safe place. Protect from light.

**CUSTOMER INFORMATION LINE**  
**1800 678 368**  
 from anywhere in Australia  
 9am to 4pm (EST) Monday to Friday  
[www.growsolutions.com.au](http://www.growsolutions.com.au)

Batch No.:   
 Expiry date:



9 510 160 824 714 >

Bayer Australia Ltd  
 875 Pacific Highway  
 Pymble NSW 2073  
 ACN 000 138 714

Manufactured in Australia.  
 VIPER® is a registered trademark of the Bayer Group.  
 APVMA Approval No. 69187/108903

20063 05/18



**POUR-ON LOUSICIDE FOR SHEEP**

**ACTIVE CONSTITUENT: 10g/L THIACTOPRID**

For the control of neotectoid susceptible and IGR resistant strains of body lice (*Bovicola ovis*) on shorn sheep up to 7 days off shears. This product can be expected to protect sheep against re-infestation with lice for 4 weeks after treatment when applied within 24 hours of shearing.

**DIRECTIONS FOR USE**

**RESTRAINTS: DO NOT USE** more than 7 days after shearing. **DO NOT USE** on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

**SHAKE WELL BEFORE USE**

**DOSAGE AND ADMINISTRATION**  
**Critical comments:** Use only on sheep that have been cleanly shorn. Avoid treating ewes less than 6 weeks before lambing commences, while ewes are lambing or have lambs at foot, as live lice can persist and infect lambs. Treated sheep should not be mixed with untreated sheep until 6 weeks after treatment. Apply the entire dose to sheep under 30kg as a single broadband stripe from poll to tail butt. For sheep over 30kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the butt of the tail. The stripes should be no further than 10cm apart. Refer to the diagrams below.

Bodyweight (kg)	Dose (mL)	Method	Apply in the pattern below
5.0 – 6.0	8		
6.1 – 8.0	12		
8.1 – 10.0	15	Single stripe	
10.1 – 12.5	20		
12.6 – 15	25		
15.1 – 20	30		
20.1 – 30	40		
30.1 – 55	60		
55.1 – 80	80	Double stripe	
80.1 – 100	90		
Add 10mL for each 10kg above 100kg.			

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Do not underdose.

Where there is a large variation in size within the group (e.g. lambs), dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**GENERAL INSTRUCTIONS**

Apply product only with the Bayer 6-stripe applicator. Sheep must be treated within 7 days of shearing. Efficacy is not affected by moderate rainfall either before or after treatment (50mm rain 1 hour prior to treatment and/or 12.5mm 2 hours after treatment and/or 2.5mm 24 hours after treatment). Treated sheep must not be mixed with lousy sheep. This product should not be used on sheep which have not been cleanly shorn for any reason eg. mycotic dermatitis.

Management practices which promote good lice control such as complete musters, accurate treatment of every animal according to label, ensuring all animals can be treated at the same time, no split shearings, stock-proof fencing to prevent external introductions, quarantine and lice treatment of all newly purchased stock are essential for effective lice control.

VIPER® Pour-On may be used in conjunction with worm control drenches and at the same time as vaccination.

VIPER® Pour-On contains a blue dye to identify treated sheep. This dye is scorable from the wool.

VIPER® Pour-On is safe for use on breeding ewes and rams.

**WITHOLDING PERIODS**

**MEAT: DO NOT USE less than 42 days before slaughter for human consumption.**  
**MILK: DO NOT USE on ewes which are producing, or may in the future produce, milk or milk products for human consumption.**  
**WOOL: DO NOT USE less than 6 months before shearing or fibre collection.**

**TRADE ADVICE**

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 56 days before slaughter for export.** Before using this product, confirm the current ESI from Bayer on 1800 678 368 or APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)).

**SAFETY DIRECTIONS**

Will damage eyes. Avoid contact with eyes. When using the product, wear cotton overalls over normal clothing, buttoned to the neck and wrist and a washable hat, PVC or rubber apron and chemical-resistant gloves and water resistant footwear and face shield or goggles. If product eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**Sheep Rehoming Interval: Four weeks. If sheep must be handled during this interval, cotton overalls buttoned to the neck and wrist (or equivalent clothing) should be worn.**

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

Additional information is listed in the Safety Data Sheet.

**GHS STATEMENTS**

**May cause an allergic skin reaction. Suspected of causing cancer if swallowed in large amounts persistently over an extended period.**  
**Prevention:** P201 Obtain special instructions before use. P261 Avoid breathing vapours/spray. P280 Wear protective gloves, protective clothing, eye protection, face protection. Response: IF ON SKIN: Wash with plenty of soap and water. P333 + P313 If skin irritation or rash occurs: Get medical advice/attention. P308 + P311 If exposed or concerned: Call a POISON CENTER/doctor/physician. Storage: P405 Store locked up. Disposal: P501 Dispose of contents/container to an approved waste disposal plant.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND**

**THE ENVIRONMENT**

Very toxic to aquatic life. DO NOT contaminate streams, rivers, wetlands or watercourses with this product or used containers.

**LIMITATION OF LIABILITY**

Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Bayer Australia Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

**DISPOSAL**

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. DO NOT dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at


any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

**STORAGE**

Store below 30°C (room temperature). Store in original container, tightly closed in a safe place. Protect from light.

**DRUM TAP ENCLOSED**

Fit to internal thread after removing small drum cap



Cut along dotted line



Bayer Australia Ltd  
 875 Pacific Highway  
 Pymble NSW 2073  
 ACN 000 138 714

Manufactured in Australia.

VIPER® is a registered trademark of the Bayer Group.

APVMA Approval No. 69187/108903

**CUSTOMER INFORMATION LINE**  
**1800 678 368**  
 from anywhere in Australia  
 9am to 4pm (EST) Monday to Friday  
[www.growsolutions.com.au](http://www.growsolutions.com.au)

Glue

Glue

250mm (height)

250mm (height)

No Image In This Area

No Image In This Area