



48.5

Selovin™ LA
Long-acting Injectable Selenium Supplement

DANGER
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



500mL

Selovin™ LA
Long-acting Injectable Selenium Supplement

For the prevention and treatment of selenium deficiency, ill thrift, white muscle disease, and all other selenium responsive diseases in sheep and cattle
Active Constituent: Each mL contains Barium selenate 177.5mg, supplying elemental **SELENIUM** 50mg.

Elanco



Elanco

Selovin™ LA
Long-acting Injectable Selenium Supplement

READ ENTIRE LABEL BEFORE USE. May be harmful if swallowed. May be harmful if absorbed through the skin. May cause eye irritation. Suspected of causing heritable mutations/genetic defects. Presumed to/may cause organ damage from repeated oral exposure at high doses. **HANDLING PRECAUTIONS** Wash hands and exposed skin before meals and after work. Avoid skin contact. Avoid contact with eyes. When handling, wear eye protection and waterproof gloves. Do not eat, drink or smoke when using this product. **FIRST AID** If swallowed, DO NOT induce vomiting. If splashed in the eyes, wash out immediately with water. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If self-injected, seek medical advice. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor, immediately. **ENVIRONMENTAL PROTECTION** Toxic to aquatic organisms with long lasting effects. Toxic to the soil environment. Avoid contamination of any water supply with product or empty container. **MODE OF ACTION** This formulation provides a rapid rise in the selenium status of sheep and cattle, reaching adequate levels in the body in 3-4 weeks and lasting up to 12 months. The treatment of animals in early to mid-pregnancy provides selenium for the newborn.

CONTRAINDICATION Do not use unless a deficiency of selenium has been diagnosed. Do not use in animals previously vaccinated with PregSure BVJ as this may lead to an increased risk of anaphylaxis. A thorough vaccination history must be obtained for the animals, both existing and new to the herd, before treatment with Barium Selenate. **DIRECTIONS FOR USE** By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label. Shake the flexipack thoroughly before and during use. Administer ONLY by subcutaneous injection into the anterior half of the neck. For requirement and frequency of dose, seek advice from your veterinarian. **SHEEP AND CATTLE** 1 mL per 50kg bodyweight Equivalent to 1mg of elemental Selenium per kg bodyweight **Lambs at weaning** 0.5mL **Calves** 1-2mL **Cows** 1-2mL **Yearling cattle** 3-5mL **Adult cattle** 6-10mL **WARNINGS** • Do not exceed dose and frequency. Do not use at the same time as any other selenium fertiliser, pill or product without consulting your veterinarian. • Do not use at the same time as any other selenium fertiliser, pill or product without consulting your veterinarian.

• In some cases, a lesion may appear at the injection site time but may last up to 3 months or longer. An intramuscular injection may cause a more serious reaction and greater care must be taken to ensure this does not occur. • Monitor the animals for the first 10-15 minutes after administration, for any hypersensitivity reactions such as increased respiration and increased heart rate, anxiety, pacing and sweating. Other signs such as swelling of the mucous membranes may be seen in some cases. Antihistamines have proven to be successful in a number of cases of hypersensitivity reactions or anaphylaxis of this sort, so this treatment should be initiated in the event of a reaction. **PRECAUTIONS** • Take care to avoid self-inoculation as this can cause serious local reaction. **It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.** **WITHHOLDING PERIODS** Milk and Meat: **NIL** **DISPOSAL** Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other

Registered pursuant to the ACMA Act 1997, No A6509
See www.elanco.co.nz for registration conditions.
Elanco New Zealand
106 Wm Station Road, Manukau, Auckland 2104
Customer Info Line: 0800 446 121

Approved facility. Crush or puncture empty container and bury in a suitable landfill. Do not use container for any other purpose.

time but may last up to 3 months or longer. An intramuscular injection may cause a more serious reaction and greater care must be taken to ensure this does not occur.

• Monitor the animals for the first 10-15 minutes after administration, for any hypersensitivity reactions such as increased respiration and increased heart rate, anxiety, pacing and sweating. Other signs such as swelling of the mucous membranes may be seen in some cases. Antihistamines have proven to be successful in a number of cases of hypersensitivity reactions or anaphylaxis of this sort, so this treatment should be initiated in the event of a reaction.

PRECAUTIONS

- Take care to avoid self-inoculation as this can cause serious local reaction.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS Milk and Meat: **NIL**

DISPOSAL Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other

consulting your veterinarian.

• Do not exceed dose and frequency. Do not use at the same time as any other selenium fertiliser, pill or product without consulting your veterinarian.



BATCH / EXPIRY

314.5

15

100

50

100

49.5

25

38

295

70 mm

100 mm

DANGER
KEEP OUT OF REACH OF CHILDREN
 FOR VETERINARY USE ONLY

Selovin LA

Long-acting Injectable Selenium Supplement

For the prevention and treatment of selenium deficiency, ill thrift, white muscle disease, and all other selenium responsive diseases in sheep and cattle

ACTIVE CONSTITUENT: Each mL contains Barium selenate 177.5mg, supplying elemental SELENIUM 50mg

DIRECTIONS FOR USE Read package insert prior to use.
By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Shake the flexpack thoroughly before and during use. Administer **ONLY** by subcutaneous injection into the anterior half of the neck. Do not exceed stated dose and frequency.

Sheep and cattle 1mL per 50kg bodyweight
 Equivalent to 1mg of elemental selenium per kg bodyweight.

Lambs at weaning	0.5mL	Yearling cattle	3 – 5mL
Sheep	1 – 2mL	Adult cattle	6 – 10mL
Calves	1 – 2mL		

PRECAUTIONS Take care to avoid self-inoculation as this can cause serious local reaction. It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHOLDING PERIODS Milk and Meat: NIL

STORAGE Store below 25°C. Protect from light. Broached packs should be discarded after 28 days.

DISPOSAL Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Crush or puncture empty container and bury in a suitable landfill.
 Do not use container for any other purpose.

Registered pursuant to the AOM Act 1997, No A9628
 See www.foodsafety.govt.nz for registration conditions.
 Elanco New Zealand
 115 Wyn Stirling Road, Marsden, Auckland 2104
 Customer Info Line: 0800 448 121




CS1005/2

MCC E1761

500mL

10-63432

DATE _____ SIGNATURE _____

APPROVED WITH MODIFICATION

APPROVED WITHOUT MODIFICATION

ROLL INFO

On / Right 3
 Core 76.2
 MaxOD : 400 mm
 Labels per roll : 500



Disclaimer: Please check carefully for text, colour separation and layout. Although artwork has been checked, MCC Ireland Ltd. accept no responsibility for errors once artwork has been approved. The label will be printed as per colours indicated, however it should be noted that this proof is not Pantone correct and colours should be checked against a Pantone Colour Guide. Artwork is to be returned signed in spec box once approved, with a digital signature and date. If you do not have the facility to do this then please approve via email with artwork attached.

DANGER

KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



Selovin™ LA

Long-acting Injectable Selenium Supplement

DESCRIPTION

A sterile, white, opaque suspension for injection. Each mL contains: Barium selenate 177.5mg supplying elemental SELENIUM 50mg.

MODE OF ACTION

This formulation produces a rise in the selenium status of sheep and cattle, reaching adequate levels in the body in approximately 3 – 4 weeks and lasting up to 12 months. The treatment of animals in early to midpregnancy provides selenium for the newborn.

INDICATIONS

For the prevention and treatment of selenium deficiency, ill thrift, white muscle disease, and all other selenium responsive diseases in sheep and cattle.

CONTRAINDICATION

Do not use unless a deficiency of selenium has been diagnosed. Do not use in animals previously vaccinated with PregSure BVD as this may lead to an increased risk of anaphylaxis. A thorough vaccination history must be obtained for the animals, both existing and new to the herd, before treatment with Barium Selenate.

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Shake the pack thoroughly before and during use. Administer **ONLY** by subcutaneous injection into the anterior half of the neck. For requirement and frequency of dose, seek advice from your veterinarian.

SHEEP AND CATTLE

1mL per 50kg bodyweight
Equivalent to 1mg of elemental Selenium per kg bodyweight

Lambs at weaning	0.5mL
Sheep	1 – 2mL
Calves	1 – 2mL
Yearling cattle	3 – 5mL
Adult cattle	6 – 10mL

WARNINGS

- Do not exceed dose and frequency. Do not use at the same time as any other selenised fertiliser, prill or product without consulting your veterinarian.
- In some cases, a lesion may appear at the injection site about a week after injection. This usually diminishes over time but may last up to 3 months or longer. An intramuscular injection may cause a more serious reaction and greater care must be taken to ensure this does not occur.

- Monitor the animals for the first 10 – 15 minutes after administration, for any hypersensitivity reactions such as increased respiration and increased heart rate, anxiety, pacing and sweating. Other signs such as swelling of the mucous membranes may be seen in some cases. Antihistamines have proven to be successful in a number of cases of hypersensitivity reactions or anaphylaxis of this sort, so this treatment should be initiated in the event of a reaction

PRECAUTIONS

- Take care to avoid self-inoculation as this can cause serious local reaction.

READ ENTIRE LABEL BEFORE USE.

May be harmful if swallowed. May be harmful if absorbed through the skin. May cause eye irritation. Suspected of causing heritable mutations/genetic defects. Presumed to/may cause organ damage from repeated oral exposure at high doses.

HANDLING PRECAUTIONS

Wash hands and exposed skin before meals and after work. Avoid skin contact. Avoid contact with eyes. When handling, wear eye protection and waterproof gloves. Do not eat, drink or smoke when using this product.

FIRST AID

If swallowed, DO NOT induce vomiting. If splashed in the eyes, wash out immediately with water. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If self-injected, seek medical advice. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor, immediately.

ENVIRONMENTAL PROTECTION

Toxic to aquatic organisms with long lasting effects. Toxic to the soil environment. Avoid contamination of any water supply with product or empty container.

WITHHOLDING PERIODS

Milk and Meat: Nil

STORAGE

Store below 25°C. Protect from light. Broached packs should be discarded after 28 days.

DISPOSAL

Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Crush or puncture empty container and bury in a suitable landfill. Do not use container for any other purpose.

Registered pursuant to the ACVM Act 1997, No A9509

See www.foodsafety.govt.nz for registration conditions.

Elanco New Zealand,
106 Wiri Station Road,
Manukau, Auckland 2104
Customer Info Line: 0800 446 121

Selovin, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.
© 2021 Elanco or its affiliates

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds