



Product Name: FASIMEC CATTLE ORAL FLUKICIDE AND BROAD SPECTRUM DRENCH  
 APVMA Approval No: 52406/134847



Label Name:	FASIMEC CATTLE ORAL FLUKICIDE AND BROAD SPECTRUM DRENCH
-------------	---

Signal Headings:	<p>CAUTION</p> <p>KEEP OUT OF REACH OF CHILDREN</p> <p>FOR ANIMAL TREATMENT ONLY</p> <p>READ SAFETY DIRECTIONS BEFORE OPENING OR USING</p>
------------------	--

Constituent Statements:	<p>120 g/L TRICLABENDAZOLE</p> <p>2 g/L IVERMECTIN</p>
-------------------------	--

Claims:	<p>Fasimec Cattle contains triclabendazole, a member of the benzimidazole family of drenches, and ivermectin, a member of the avermectin (macrocyclic lactone) family of drenches.</p> <p>For the treatment of triclabendazole-sensitive Early Immature, Immature and Mature liver fluke (<i>Fasciola hepatica</i>) in cattle.</p> <p>For the treatment and control of ivermectin-sensitive strains of roundworms and lungworm (including benzimidazole, levamisole, and morantel-resistant strains), and sucking lice in cattle.</p> <p>Parasites treated and controlled by Fasimec Cattle        Early immature, immature and adult liver fluke <i>Fasciola hepatica</i></p> <p>Adult and immature (L4) stages of:        Small brown stomach worm (including inhibited stages) <i>Ostertagia ostertagi</i>        Barber's pole worm <i>Haemonchus placei</i>        Stomach hair worm <i>Trichostrongylus axei</i>        Intestinal hair worm <i>Trichostrongylus colubriformis</i>        Small intestinal worm <i>Cooperia</i> spp.        Nodule worm <i>Oesophagostomum radiatum</i></p> <p>Adult stages of:        Large bowel worm <i>Oesophagostomum venulosum</i>        Whipworm <i>Trichuris</i> spp.        Intestinal threadworm <i>Strongyloides papillosus</i></p> <p>Immature stages of:        Thin necked intestinal worm <i>Nematodirus</i> spp.</p> <p>Adult and immature stages of:</p>
---------	--

	<p>Lungworm Dictyocaulus viviparus</p> <p>Sucking lice:  Long-nosed sucking louse Linognathus vituli  Short-nosed sucking louse Haematopinus eurysternus  Small blue sucking louse Solenopotes capillatus</p> <p>Does not provide control of the cattle biting louse, Bovicola (Damalinia) bovis.</p> <p>Resistance may develop to any drench. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a drench resistance test be conducted before any parasite treatment is used.</p>
--	---

Net Contents:	1 L 2.5 L 5 L 5 L Gunpack 15 L 20 L
---------------	--

Directions for Use:	
---------------------	--

Restrains:	NOT TO BE USED FOR ANIMALS PRODUCING MILK FOR HUMAN CONSUMPTION OR PROCESSING.
------------	--

Contraindications:	
--------------------	--

Precautions:	In areas where Black Disease is known to occur, adequate vaccination procedures are advised.
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	<p>SHAKE WELL BEFORE USE</p> <p>ORAL DOSE RATE: CATTLE 5 mL per 50 kg BW</p> <p>Critical Comments:</p> <ol style="list-style-type: none"> <li>1. FASIMEC may be administered at the same time as animals are treated for biting lice.</li> <li>2. FASIMEC may be administered to calves, pregnant cows or weak or debilitated animals.</li> <li>3. Wash out equipment with water after use.</li> <li>4. FASIMEC is palatable to cattle.</li> <li>5. Cattle in excess of 650 kg should be dosed at 5 mL per 50 kg. A representative sample of animals should be weighed before treatment either with scales or a weigh band. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). DO NOT underdose. Where there is a larger variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.</li> </ol> <p>This section contains file attachment.</p>
----------------------------	---

General Directions:	<p>FASIMEC Cattle is a ready to use, free flowing aqueous suspension for the treatment and control of acute and chronic fascioliasis, mixed infections of gastrointestinal roundworms, lungworm and sucking lice.</p> <p>Fasimec Cattle when used at the recommended rate of 12 mg triclabendazole and 0.2 mg ivermectin per kg body weight for cattle, i.e., 5 mL Fasimec Cattle per 50 kg body weight, combines high efficacy against all ages of triclabendazole-sensitive liver fluke, ivermectin-sensitive adult and immature roundworms, lungworm and sucking lice, with a high safety margin.</p> <p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p>
Withholding Periods:	<p>WITHHOLDING PERIODS</p> <p>MEAT: DO NOT USE less than 21 days before slaughter for human consumption.</p> <p>MILK: DO NOT USE in lactating cows or within 28 days of calving where milk may be used or processed for human consumption.</p>
Trade Advice:	<p>TRADE ADVICE–EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 56 days before slaughter for export. Before using this product, confirm the current ESI from the Elanco Australasia Pty Ltd on 1800 226 324 or the APVMA website (<a href="http://www.apvma.gov.au/residues">www.apvma.gov.au/residues</a>).</p>
Safety Directions:	<p>May irritate the eyes. Avoid contact with eyes. Wash hands after use.</p>
First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.</p>
First Aid Warnings:	
Additional User Safety:	<p>If additional hazard information is required refer to the Safety Data Sheet. For a copy phone 1800 633 768.</p>
Environmental Statements:	<p>ENVIRONMENTAL PROTECTION:</p> <p>Protection of Wildlife, Fish, Crustaceans and the Environment: Ivermectin is extremely toxic to aquatic species. DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used container.</p>
Disposal:	<p>1 L container:</p> <p>Dispose of container by wrapping with paper and putting in garbage.</p> <p>2.5 L, 5 L, 15 L and 20 L containers:</p> <p>Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If the container has the DrumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any DrumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management</p>

	<p>facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
--	---

Storage:	<p>Store below 30 °C (room temperature) in original container, tightly closed, in a cool, dry place. Protect from light.</p>
----------	--

**DOSAGE AND ADMINISTRATION: ORAL DOSE RATE: CATTLE 5 mL per 50 kg BW**

<b>Parasites controlled</b>	<b>Cattle weight (kg)</b>	<b>Oral dose (mL)</b>
Liver fluke, Gastrointestinal roundworms, Lungworm and Sucking Lice	Up to 100	1 mL/10 kg body weight
	101–150	15
	151–200	20
	201–250	25
	251–300	30
	301–400	40
	401–500	50
	501–600	60
	601–650	65
	Over 650	5 mL/50 kg body weight