Instructions

Fill in the light blue shaded boxes.
The remaining values will be calculated from this input.



Mg monensin per head per day	
Rumensin® dose requested per head per day	mg
Amount of medicated feed to be fed per day, as fed	kg
Dry Matter Intake	kg
Body weight of cattle being fed	kg
DMI expressed as % of body weight	%
Equivalent in mg/kg (ppm) monensin in complete diet, 100% DM	mg/kg (ppm)
Monensin required per kg of medicated feed, as fed	mg/kg
Amount of Rumensin Premix to add per tonne of medicated feed, as fed	kg/tonne
Completed by:	Date (dd-mm-yyyy):

Notes:

INDICATIONS:

- 1. For improved feed efficiency in beef cattle (steers and heifers) fed in confinement for slaughter (33 to 48 mg/kg).*
- 2. As an aid in the prevention of coccidiosis caused by Eimeria bovis and Eimeria zuernii in cattle (22 mg/kg).
- 3. For increased rate of weight gain in growing cattle on pasture (slaughter, stocker and feeder cattle, and beef and dairy replacement heifers) of greater than 180 kg (400 lb) body weight (200 mg per head per day).

- mg per nead per day).

 4. For reduction of milk fat percentage in lactating dairy cows (16 to 24 mg/kg).*

 5. For minimizing loss of body condition during lactation in dairy cows (8 to 24 mg/kg).*

 6. For improving feed efficiency of milk protein production in lactating dairy cows (16 to 24 mg/kg).*
- 7. For the prevention of coccidiosis associated with pathogenic *Eimeria spp.* in sheep (11 to 22 mg/kg)*
 8. For the prevention of coccidiosis associated with *Eimeria crandallis*, *E. christenseni*, and *E. ninakohlyakimovae* in goats (11 to 22 mg/kg)* Indications in mg/kg (ppm) are calculated for complete diets corrected to a 100% dry matter basis.

For all claims except #3: Thoroughly mix supplements and premixes in the total daily diet or in complete feed (grain portion of the ration) before use. Do not feed undiluted. For claim #3: The medicated supplement must be prepared to be hand fed daily at a minimum of 0.5 kg/head/day to deliver 200 mg of monensin/ head/day. The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

Consumption by unapproved species or feeding undiluted may be toxic or fatal.

Rumensin, Elanco and the diagonal bar are trademarks of Elanco.

© 2025 Elanco or its affiliates. PM-CA-25-0309

