





EAN 13 Barcode
 100% Magnification
 BATCH / EXPIRY

Distributed by:
Elanco New Zealand
 106 Wiri Station Road, Manukau, Auckland 2104.
 Customer Info Line: 0800 446 121
 Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates © 2023 Elanco or its affiliates.
 New Zealand patent application no. 802089

HSN0 No. HSR002521
 Registered in New Zealand by:
Biocell Corporation Ltd.
 19 Laureston Ave, Manukau, Auckland.
 New Zealand. Phone: 09 2700755

Registered pursuant to the AVM Act 1997.
 No A011894.
 See www.foodsafety.govt.nz for registration conditions.

WITHHOLDING PERIODS
 No withholding period is required when used as directed.
Additional: Every 2 months in wet conditions
 3 - 4 weeks before mating
Heifers: Every 3 months, especially 3 - 4 weeks before weaning
Calves: At day 1 and weaning
Cows: 3 - 4 weeks before calving, mating and drying off
Cattle: over 2 years 1mL/100kg
Calves: 1 - 2 years 1 mL/75 kg
Calves: up to 1 year 1mL/50 kg

PRECAUTIONS:
 Not to be used concurrently with any other form of copper supplementation. Do not use at the same time as any other sensitised fertiliser, pill or product and do not exceed the stated dose or frequency without consulting a veterinarian. Do not administer to severely debilitated animals. Do not administer to animals suffering from liver disease, fascioliasis or which have been grazing on plants which may cause liver disease. Do not use at the same time as any other sensitised fertiliser, pill, or product without consulting a veterinarian. Do not exceed the stated dose. Do not administer to emaciated animals of body condition score less than 3/10.

CONTRAINDICATIONS:
 Not to be used concurrently with any other form of copper supplementation. Do not administer to animals suffering from liver disease, fascioliasis or which have been grazing on plants which may cause liver disease. Do not use at the same time as any other sensitised fertiliser, pill, or product without consulting a veterinarian. Do not exceed the stated dose. Do not administer to emaciated animals of body condition score less than 3/10.

ACTIVE CONSTITUENTS
 Zinc 40 mg/ml (as disodium zinc EDTA), manganese 10 mg/ml (as disodium manganese EDTA), copper 15 mg/ml (as copper disodium EDTA), selenium 5 mg/ml (as sodium selenite) and 1.4 mg/ml Vitamin B₁₂ (cyanocobalamin).

DIRECTIONS FOR USE
 By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label. For use in cattle by subcutaneous injection.

READ ENTIRE CARTON BEFORE USE.
 Only treat animals in good health. Minimise stress to animals during handling for administration.
 Slight local reaction may occur for about 30 seconds after injection. A slight swelling may be observed at the injection site for a few days after administration. Use standard aseptic procedures during administration of injections. Doses greater than 7 ml should be administered in separate sites. Inject MULTIBOOST WITH B₁₂ on the opposite side of the neck to any other products or any other injection sites.
DOSAGE RECOMMENDATIONS
 Cattle: 1 - 2 years 1 mL/75 kg
 Calves: up to 1 year 1mL/50 kg

500mL

STORAGE
 Store below 25 °C in original box in a dry place. Protect from light. Keep container well closed. Following withdrawal of the first dose, unused product should be discarded after 9 months.
 Harmful to aquatic life with long lasting effects. Toxic to terrestrial vertebrates. Avoid release to the environment. Collect spillage. Store locked up. at hand.

IF SWALLOWED: Immediately call a Poison Information centre (0800 POISON (0800 764 766) or a doctor. Rinse mouth. Causes mild skin irritation and serious eye irritation, if skin or eye contact occurs. Get medical advice. Wear eye protection.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. May cause damage to the liver through prolonged or repeated exposure. Do not breathe spray/mist. Get medical advice if you feel unwell. If medical advice is needed, have product container or label at hand.

Chelated trace element injection for beef and dairy cattle containing zinc, manganese, selenium, copper and vitamin B₁₂ for supplementation.

Injectable Trace Minerals with Vitamin B₁₂ for Cattle
WITH B₁₂
MULTIBOOST

Elanco

KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY
MULTIBOOST B₁₂
 Injectable Trace Minerals with Vitamin B₁₂ for Cattle

Elanco

500mL

KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY

Elanco

MULTIBOOST WITH B₁₂

Injectable Trace Minerals with Vitamin B₁₂ for Cattle

Chelated trace element injection for beef and dairy cattle containing zinc, manganese, selenium, copper and vitamin B₁₂ for supplementation.

Zn Mn Se Cu B₁₂

CONTRAINDICATIONS:
 Not to be used concurrently with any other form of copper supplementation. Do not administer to animals suffering from liver disease, fascioliasis or which have been grazing on plants which may cause liver disease. Do not use at the same time as any other sensitised fertiliser, pill, or product without consulting a veterinarian. Do not exceed the stated dose. Do not administer to emaciated animals of body condition score less than 3/10.

500mL