

Fusogard® vaccine is the simple and straightforward way to decrease the severity of footrot and keep your breeding program on track.

The major cause of footrot is bacterium *Fusobacterium necrophorum*, an organism commonly found in the digestive tract of cattle.

Any surface contaminated with manure may contain this organism. *F. necrophorum* can also escape from the rumen and circulate internally. An animal's environment plays a role in the development of footrot. Wet, muddy conditions or rough, abrasive terrain can cut or bruise the skin between the toes, creating the right conditions for the bacteria to invade, multiply and cause disease.

<b>FUSOGARD®</b>	Fusobacterium Necrophorum Bacterin For Use In Animals Only	
FOOTROT	INDICATIONS	For use in healthy cattle 6 months of age or older as an aid in the reduction of clinical signs of footrot caused by <i>Fusiformis necrophorus</i> ( <i>Fusobacterium necrophorum</i> ).
	ADJUVANT	SuprImm™.
	DIRECTIONS	Shake well before using. Administer 2 mL subcutaneously. In accordance with Beef Quality Assurance guidelines, this product should be administered subcutaneously (under the skin) in the neck. Revaccinate in 3 weeks. Revaccinate annually or as recommended by your veterinarian.
	PRECAUTIONS	Store out of direct sunlight at 2°-8°C (35°-46°F). DO NOT FREEZE. Use entire contents when first opened. Do not mix with other products, except as specified on the label. In case of human exposure, contact a physician. Do not vaccinate within 21 days prior to slaughter. Anaphylactic reactions may occur. Symptomatic treatment: Epinephrine.

The label contains complete use information, including cautions and warnings. Always read, understand and follow the label and use directions.

Fusogard, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.  
 ©2021 Elanco PM-CA-21-0535

