
Top



130 mm

For a copy please go to www.elanco.co.nz.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds. WITHHOLDING PERIODS MEAT: When applied to sheep off-shears or with up to 3 months wool: NIL MEAT: When applied to sheep with at least 6 months wool (Emergency treatment): Sheep producing meat or offal for human consumption must not be sold for slaughter for 7 days following treatment. WOOL: NIL MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment. Storage: Store below 30 °C in a dry place in a tightly closed container. Do not freeze. Do not store in direct sunlight. Registered to: Elanco Australasia Pty Ltd 27/88 Shortland Street Auckland, 1010 New Zealand HELPLINE: Phone 0800 426 633 www.elanco.co.nz Registered pursuant to the ACVM Act 1997, No A010205 See www.foodsafety.govt.nz for registration conditions Approved pursuant to the HSNO Act 1996, No HSR100008 See www.epa.govt.nz for approval conditions For advice on your dipping program, contact your supplier or call the Elanco Helpline: 0800 426 633. Extinosad Pour-On, Elanco and the diagonal bar logo are trademarks of

on each side of the midline from the poll to the base of the tail). GENERAL INFORMATION Optimum lice control requires the application of good agricultural practice. This includes ensuring a complete muster so that no animals remain untreated, attention to stock security following treatment, quarantine treatment of new stock arrivals, thorough and accurate dipping management and utilising the principles of Integrated Pest Management recommended by Wool Pro. DO NOT mix treated sheep with untreated sheep. It is good agricultural practice not to treat animals when they are wet or if rain is likely. However moderate rainfall following treatment does not adversely affect the effectiveness of the product. Prolonged heavy rain following treatment may reduce efficacy. ExPO contains a blue dye to identify treated sheep. The dye is scourable from the wool. Do not leave container, hose or applicator in the sun for long periods of time. At the end of use, clean out the applicator, nozzle and hose by pumping through with clean water. 107 mm DIRECTIONS FOR USE: For the control of lice on all breeds of sheep. SHAKE CONTAINER GENTLY SEVERAL TIMES BEFORE USE By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed Use any standard T-bar pour-on nozzle, and an applicator with a total adjustable volume ideally no greater than 20 mL. Do not under dose – check the accuracy of equipment before and during treatment. Coarse wool breeds: Apply ExPO Extinosad Pour-On evenly along the backline from the poll to the base of the tail using a suitable applicator with a T-bar nozzle. In sheep weighing less than 35 kg apply

as one band and in sheep weighing more than 35 kg, halve the dose volume and apply in

2 bands, approximately 5 cm either side of the midline of the back. This application pattern

ExPO can be used on up to 3 month's wool growth, but for optimum results treatment is

Body weight Dose This pack will treat

10

16.5

12.5

15

(kg) (mL) (number of sheep)

>75 Add 1 mL for each 4.5 kg above 75 kg

>75 Add 1 mL for each 4 kg above 75 kg

Apply ExPO Extinosad Pour-On evenly along the backline from the poll to the base of the tail using a suitable applicator with a T-bar nozzle. In sheep weighing less than 25 kg apply as one band and in sheep weighing more than 25 kg, halve the dose volume and apply in

1000

666

606

800

Dosage and administration coarse woolled breeds of sheep

Book Overlay 132 x 100 mm

Base

Dose

(kg)

20-25

26-30

31-35

36-40

41-50

51-60

61-70

132

mm

on the label.

will improve the spread of the product.

<35

35-55

55-75

<35

35-55 17.5

3 months after 55-75 20 500

2 bands, approximately 5 cm either side of the midline of the back.

recommended off-shears.

Off-shears

(within 24 hours

after shearing)

Sheep from

24 hours to

shearing

Fine wool breeds:

(mL)

12.5

16

19

22

25

30

35

>70 Add 2.5 mL for each 5 kg above 70 kg

penned closely and held in the yards immediately after application.

Emergency Treatment Coarse and Fine Woolled Sheep

Dosage and administration fine woolled breeds of sheep off-shears treatment only

This pack will treat

(number of sheep)

800

625

526

454

400

333

285

ExPO Extinosad Pour-On is spread over the body by passive diffusion and physical contact

(e.g. rubbing with other sheep). To facilitate product movement, treated sheep should be

Apply a 50 mL dose to reduce louse numbers in severely infested long woolled sheep

(>6 months after shearing). This dose should be applied as 4 x 12.5 mL bands (two bands



100 mm

Booklet pages must be in multiples of 4. You may delete or add pages in multiples of 4 to fit artwork.