	Full Dose (6 weeks)	Top Up Dose (4 weeks)
251 – 330kg	3 boluses	2 boluses
331 – 440kg	4 boluses	2 boluses
441 – 550kg	5 boluses	3 boluses
551 – 660kg	6 boluses	3 boluses

WARNING
Slight resistance may be felt during the early stages of the positioning of the applicator as the animal uses its tongue to try and prevent the bolus entering the back of the mouth. Always wait until the swallowing reflex is operating before administering the boluses. The swallowing reflex can also be used to assist in the correct positioning of the applicator in the mouth. Administration when the head and neck are at the wrong angle, or the use of excess force, can damage the back of the animal's throat or lead to accidental application into the windpipe, which can result in serious consequences, some of which may be fatal.

CONTRAINDICATIONS
Do not use in conjunction with other forms of zinc therapy.
Do not use at times other than the facial eczema risk period.
Do not administer with any other intra-ruminal boluses.
Administration of zinc may depress tissue copper levels in areas of copper deficiency.

WITHOLDING PERIOD
Meat: Nil
Milk: Nil

STORAGE
Store below 25°C (room temperature) in a dry place.
Store away from direct sunlight. Do not freeze.

DISPOSAL
Preferably dispose of the product by use. Otherwise dispose of product and packaging in an approved landfill or other approved facility. Crush or puncture and bury in a suitable landfill, or if appropriate recycle.

READ ENTIRE LABEL BEFORE USE.
FIRST AID:
For advice, contact the National Poisons Centre:
0800 POISON (0800 764 766) or a doctor.

Registered pursuant to the ACVM act 1997, No. A10744.
See www.foodsafety.govt.nz for registration conditions.

ELANCO NEW ZEALAND
106 Wiri Station Road, Manukau, Auckland 2104.
www.elanco.com | 0800 446 121.

5 420036 978961

Face-Guard, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.
© 2023 Elanco or its affiliates

BATCH / EXPIRY

SH504949X

INDICATIONS
Face-Guard 251 – 660kg is indicated for the prevention of facial eczema in cattle for up to 6 weeks following one treatment, or for 10 weeks if a top up dose is given six weeks after the first dose.

DIRECTIONS FOR USE FOR ORAL USE
By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Presentation
The product packaging contains the boluses conveniently packed in pairs taped together by paper wrapper. After administration the wrapper dissolves in the rumen, allowing the two boluses to separate.

First Dose
Using the applicator administer boluses at the beginning of the facial eczema risk period, when spore counts begin to rise, (mid-January and February in most areas). **This dose will provide up to 6 weeks' protection.**

Cattle:
- **251 – 330kg bwt = 3 x 132g boluses**
- **331 – 440kg bwt = 4 x 132g boluses**
- **441 – 550kg bwt = 5 x 132g boluses**
- **551 – 660kg bwt = 6 x 132g boluses**

Top-up dosing
Where there is a prolonged risk of facial eczema, a top up dose can be given after 6 weeks to provide at least 4 weeks' additional protection. **A top-up dose is 2 x 132g boluses for all cattle between 251kg and 440kg, or 3 x 132g boluses for all cattle between 441kg and 660kg.**
Thereafter, further treatment intervals should be based on faecal zinc concentrations, the level of the facial eczema risk, and veterinary advice.

FGI132-23/2

General Dosing Instructions
To ensure the safe administration of the boluses follow the steps listed:

Cattle Restraint
The animal should ideally be restrained in a cattle crush.
Avoid severe restraint, such as gripping the tongue, raising the nose too high or the use of bulldog nose clamps, as this interferes with swallowing.

Correct positioning of the Head and Neck
Administer the boluses slowly and with care, making sure the animal's head and neck are extended and straightened so the applicator in the mouth lines up with the opening in the throat.

Administering the boluses
Place the boluses in the special applicator provided with the rounded ends facing outwards. 2, 3, or 4 boluses can be administered in a single application. Where 5 or 6 boluses are to be given, these should be administered in two applications, one involving the dosing of 3 (for a total dose of 5) or 4 boluses (for a total dose of 6) and the other involving the dosing of the remaining 2 boluses.

The boluses held within the applicator should then be gently introduced into the side of the mouth and over the tongue. When positioned correctly, advance the applicator plunger with steady gentle pressure once the animal begins to swallow. Only continue administering the boluses while the animal is swallowing. If the head and neck are at the wrong angle the animal will stop swallowing and administration must cease. Administration must only recommence when the head and neck have been repositioned correctly.

Regurgitation of the bolus can occur following administration, but experience has shown that the incidence is usually of a low order. Unless severely damaged (ie. broken into two or more pieces or with significant amount of bolus missing), regurgitated boluses can be re-administered. Watch the animal for a few moments following removal of the applicator.

PRODUCT INFO

BLDE #	SH504949X	Product Name:	Face-Guard Bolus 251-660kg
Base Code:	SH504949X	Component:	Carton
Product Code:	18444	Pack Size:	251-660kg
Previous Item Code:	SH311135C		

ARTWORK INFO

Template:	133 x 89 x 106mm	Packaging Specific:	N/A
Barcodes Type:	EAN 13: 5420036978961	Add. Info:	N/A
GTIN:	GTIN Not Required	Minimum Core Dots Point Size:	6.5pt
		Proof #	JS 21-APR-2023
		Pre-Flight Date:	

NON PRINTING	CYAN	MAGENTA	YELLOW
BLACK			

ELANCO ARTWORK LEGEND v16