

205mm

90mm

**READ ENTIRE LABEL BEFORE USE**

**WARNING**

Harmful if swallowed. Causes skin irritation. Causes serious eye irritation.

**HANDLING PRECAUTIONS:** Wear protective gloves. Wear eye or face protection. Do not eat, drink or smoke when using this product. Wash thoroughly after handling.

**FIRST AID:** IF SWALLOWED: Call a POISON CENTER or doctor if you feel unwell. Rinse mouth. Take off contaminated clothing and wash it before reuse. IF ON SKIN: Wash with plenty of water. If skin irritation occurs: Get medical advice or attention. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice or attention. For advice contact the National Poisons Centre 0800 POISON (0800 764 766). If medical advice is needed, have the product container or label at hand.

**EMERGENCY RESPONSE:** Stop leak if without risk. Move containers from spill area. Dilute with water and mop up if water-soluble. Alternatively, or if water-insoluble, absorb with an inert dry material and place in an appropriate waste disposal container. For specialist advice in an emergency call CHEMTREC 0800 425 459. If additional hazard information is required, refer to the Safety Data Sheet. For a copy please go to [www.elanco.co.nz](http://www.elanco.co.nz).

**DISPOSAL:** Treat the product so that it is no longer toxic. Preferably dispose of the product by use. Otherwise dispose of product and packaging as hazardous waste in an approved landfill or other approved facility.



**WARNING**

KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY

**ORAL-CAL™**  
Oral Calcium Supplement

**ACTIVE CONSTITUENT:** Each mL contains 490mg CALCIUM CHLORIDE DIHYDRATE equivalent to 53.4g calcium per 400mL bottle

As an aid in prevention and treatment of milk fever (hypocalcaemia) in cows. Also as a calcium supplement where deficiency of dietary calcium may occur.



**DIRECTIONS FOR USE:** By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

**FOR ORAL USE ONLY**

Take care when drenching, especially if cow is showing signs of paralysis. Direct the neck of the bottle over the tongue towards the back of the throat and slowly squeeze the bottle to assist swallowing. Avoid accidental administration into the windpipe and lungs.

**Prevention of milk fever:** Give one bottle (400mL) 24 hours before calving, one bottle at calving, one bottle 10 – 12 hours after calving and one bottle 24 hours after calving.

**Treatment of milk fever:** To be used in conjunction with intravenous calcium treatment. Administer one bottle (400mL) per cow as soon as the cow is standing and can swallow normally. Repeat with the same dose 10 – 12 hours later.

**Calcium deficiency – when unrelated to calving time:** Give one bottle (400mL) morning and night for 2 days.

**WITHHOLDING PERIODS:** Milk and Meat: **NIL**  
**STORAGE:** Store below 25°C (room temperature).

BATCH / EXPIRY

**Elanco™**

**400mL**

YL701441A

Registered pursuant to the ACVM Act 1997, No A007669  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.  
Elanco New Zealand,  
106 Wiri Station Road, Manukau, Auckland 2104  
Customer Info Line: 0800 446 121



**PRODUCT INFO**

BLUE #:	701441APB	Product Name:	ORAL CAL
Item Code:	YL701441A	Component:	Label
Product Code:	N/A	Pack Size:	400mL
Previous Item Code	90198313		

**ARTWORK INFO**

Template:	LA_205x90mm_v1a	Packaging Spec(s):	N/A
Barcodes/Type:	EAN 13: 5420036977605	Add. Info:	N/A
		Minimum Core Data Point Size:	6 pt
GTIN:	GTIN Not Required	Proof #	P1a
		By/Date	TCK 24-OCT-2023

ELANCO ARTWORK LEGEND v16

<b>LEGEND KEY</b>	Template: LA_205x90mm_v1a
Text Free Area	
	50 mm Rule Bar

Elanco Template Key v1