


INJECTION SITE SWELLING



REDUCING INJECTION SITE SWELLING

- Proper vaccination location
- · Size and length of needle
 - 16 gauge x 1.5 inches
- Cleanliness (dry, free from mud and manure)
- Change needles often to reduce the use of burred needles
- · Proper restraint to reduce tissue damage
- Nutritional status of herd



Breeding date	Calving date	Scour Bos® 9 – Heifers/Naive females – Primary year			Scour Bos 4
			Annual revaccination subsequent calvings		Primary year only
		16 weeks	10 weeks	8 weeks	4 weeks
1-Apr	9-Jan	19-Sep	31-Oct	14-Nov	12-Dec
15-Apr	23-Jan	3-Oct	14-Nov	28-Nov	26-Dec
1-May	8-Feb	19-Oct	30-Nov	14-Dec	11-Jan
15-May	22-Feb	2-Nov	14-Dec	28-Dec	25-Jan
1-Jun	11-Mar	19-Nov	31-Dec	14-Jan	11-Feb
15-Jun	25-Mar	3-Dec	14-Jan	28-Jan	25-Feb
1-Jul	10-Apr	19-Dec	30-Jan	13-Feb	13-Mar
15-Jul	24-Apr	2-Jan	13-Feb	27-Feb	27-Mar
1-Aug	11-May	19-Jan	2-Mar	16-Mar	13-Apr
15-Aug	25-May	2-Feb	16-Mar	30-Mar	27-Apr
1-Sep	11-Jun	19-Feb	2-Apr	16-Apr	14-May
15-Sep	25-Jun	5-Mar	16-Apr	30-Apr	28-May
1-Oct	11-Jul	21-Mar	2-May	16-May	13-Jun
15-Oct	25-Jul	4-Apr	16-May	30-May	27-Jun
1-Nov	11-Aug	21-Apr	2-Jun	16-Jun	14-Jul
15-Nov	25-Aug	5-May	16-Jun	30-Jun	28-Jul
1-Dec	10-Sep	21-May	2-Jul	16-Jul	13-Aug
15-Dec	24-Sep	4-Jun	16-Jul	30-Jul	27-Aug
1-Jan	11-Oct	21-Jun	2-Aug	16-Aug	13-Sep
15-Jan	25-Oct	5-Jul	16-Aug	30-Aug	27-Sep
1-Feb	11-Nov	22-Jul	2-Sep	16-Sep	14-Oct
15-Feb	25-Nov	5-Aug	16-Sep	30-Sep	28-Oct
1-Mar	9-Dec	19-Aug	30-Sep	14-Oct	11-Nov
15-Mar	23-Dec	2-Sep	14-Oct	28-Oct	25-Nov
Make sure there is a minimum of 28 days between the 1st & 2nd vaccination the primary year, per					

Make sure there is a minimum of 28 days between the 1st & 2nd vaccination the primary year, per label indications. This ensures that the animal's immune system recognizes the 2nd dose as a booster.