



Product Name: BAYCOX COCCIDIOCIDICIDE FOR PIGLETS AND CATTLE
APVMA Approval No: 68946/130682

Label Name:	BAYCOX COCCIDIOCIDICIDE FOR PIGLETS AND CATTLE
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS FOR ANIMAL TREATMENT ONLY
Constituent Statements:	50 g/L TOLTRAZURIL
Claims:	<p>For the control of coccidiosis in neonatal piglets caused by <i>Isospora suis</i>, and in cattle up to 9 months of age caused by <i>Eimeria bovis</i> or <i>Eimeria zuernii</i>.</p> <p>BAYCOX is effective in one dose because it attacks all stages of the parasite in the animal.</p> <p>Piglets: BAYCOX controls coccidiosis caused by <i>Isospora suis</i> infection of young piglets. <i>Isospora suis</i> infection is a major cause of diarrhoea, dehydration, loss of appetite, poor growth, and occasional mortality in piglets between birth and 3-4 weeks of age.</p> <p>Cattle: BAYCOX is for the treatment of clinical coccidiosis, and as an aid in the prevention of coccidiosis in cattle up to 9 months of age. BAYCOX controls coccidiosis caused by <i>Eimeria bovis</i> or <i>Eimeria zuernii</i> in young cattle. Coccidiosis typically occurs in cattle less than a year old causing diarrhoea which is often blood stained. Recently weaned calves are susceptible to outbreaks of clinical disease.</p> <p>Resistance may develop to any chemical.</p>
Net Contents:	250 mL 1 L 1 L (backpack)
Directions for Use:	
Restrains:	DO NOT RE-TREAT calves for 75 days after the last treatment. DO NOT TREAT piglets later than 8 days of age.

Contraindications:	
Precautions:	
Side Effects:	
Dosage and Administration:	<p>SHAKE THE CONTAINER BEFORE USE</p> <p>Give by mouth. Ensure the applicator is delivering a full dose before starting treatment. The applicator should be set to deliver a full dose per stroke.</p> <p>This section contains file attachment.</p>
General Directions:	<p>Piglets: In the case of diarrhoea in neonatal piglets, obtain a veterinary diagnosis of coccidiosis before using this product. Clinical coccidiosis of piglets appears in piglets from 5 to 30 days of age. Control of the disease is most effective when BAYCOX is given as a preventive before intestinal damage occurs and diarrhoea begins.</p> <p>Cattle: In the case of diarrhoea in calves, obtain a veterinary diagnosis of coccidiosis before using this product. Early treatment of calves as soon as the diagnosis is obtained will prevent further intestinal damage in affected animals and prevent infection of the herd. To obtain maximum benefit on farms with a history of coccidiosis, BAYCOX should be given approximately 1 week before expected onset of clinical signs. Treatment of newly weaned calves at the time of milk withdrawal will control coccidiosis associated with weaning.</p>
Withholding Periods:	<p>MEAT (PIGS): DO NOT USE less than 70 days before slaughter for human consumption.</p> <p>MEAT (CATTLE): DO NOT USE less than 56 days before slaughter for human consumption.</p> <p>MILK (CATTLE): DO NOT USE in lactating or pregnant cows where milk may be used or processed for human consumption.</p>
Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI) PIGS: An ESI has not been established for this product for pigs. Note — observing the meat withholding period may not be sufficient to mitigate potential risks to export trade. Trade advice should be sought from Elanco Australasia Pty Ltd on 1800 995 709 before using this product.</p> <p>EXPORT SLAUGHTER INTERVAL (ESI) CATTLE: DO NOT slaughter cattle for export for 75 days after last treatment. The ESI for cattle on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Elanco Australasia Pty Ltd on 1800 995 709 or the APVMA website (www.apvma.gov.au/residues)</p>
Safety Directions:	When using the product wear rubber gloves. After each day's use wash gloves.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
First Aid Warnings:	
Additional User Safety:	

Environmental Statements:	ENVIRONMENTAL PROTECTION: Do not contaminate ponds, waterways or drains with this product or used container.
---------------------------	---

Disposal:	Dispose of container by wrapping with paper and putting in the garbage.
-----------	---

Storage:	Store below 30°C (room temperature)
----------	-------------------------------------

Baycox Coccidiocide for piglets and cattle

PN 68946

Dose table:

	Dose	Prevention	Treatment
Piglets	1 mL	Treat at 3-6 days old	Treat at onset of coccidial diarrhoea. Treat all piglets in infected litters and all infected and 'at risk' litters in the farrowing house.
Cattle	3 mL per 10 kg bodyweight	In the case of history of coccidiosis on farm, treat 1 week before expected onset of clinical signs. For prevention of coccidiosis in recently weaned calves, treat at weaning time when milk feeding ceases.	Treat at onset of diagnosed clinical signs