



Product Name: COFORTA 100 INJECTABLE PHOSPHORUS SOLUTION  
APVMA Approval No: 50277/130153

Label Name:	COFORTA 100 INJECTABLE PHOSPHORUS SOLUTION
-------------	--

Signal Headings:	FOR ANIMAL TREATMENT ONLY
------------------	---------------------------

Constituent Statements:	100 mg/ml BUTAPHOSPHAN 50 µg/ml VITAMIN B12 = CYANOCOBALAMIN  The solution contains 17.3 mg of phosphorus in 1 mL
-------------------------	--

Claims:	For the supplementation of animals with phosphorus where it is apparent that such treatment would be beneficial to the animal's condition and where the simultaneous administration of Vitamin B12 is required.
---------	---

Net Contents:	100mL
---------------	-------

Directions for Use:	
---------------------	--

Restraints:	
-------------	--

Contraindications:	
--------------------	--

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	Dispose of broached vials within 28 days of opening.  For I.M. or S.C. injection according to enclosed directions for use. Sterile technique and equipment must be used for each injection.  To be used only by or under the supervision of a veterinary surgeon. Give by intramuscular or subcutaneous injection according to the following table. Volumes of more than 20 mL should be split and given at several intramuscular sites or given subcutaneously.
----------------------------	---

	<p>In acutely diseased animals:</p> <p>Horse, Cattle 5 - 25 mL  Foal, Calf 5 – 12 mL  Sheep, Goat 2.5 – 5 mL  Lamb 1.5 – 2.5 mL  Pig 2.5 – 10 mL  Suckling pig 1 – 2.5 mL  Dog 0.5 – 5 mL  Cat 0.5 – 5 mL</p> <p>Repeat daily if required.</p> <p>In chronic disease give half the above dosage at intervals of 1-2 weeks or less. In healthy animals give half the above dose.</p>
--	---

General Directions:	<p>Caution: avoid carcass damage</p> <ol style="list-style-type: none"> <li>1. Sterilise all injection apparatus by boiling before use. Avoid use of strong disinfectants on apparatus.</li> <li>2. Maintain cleanliness at all times.</li> <li>3. Keep needles sharp and clean; replace frequently.</li> <li>4. Use needles of appropriate gauge and length. For subcutaneous administration, use shortest needle possible, certainly not exceeding 15 mm.</li> <li>5. As far as possible avoid injection of animals in wet weather or under dusty conditions.</li> <li>6. Intramuscular injections should be made into muscle tissue on the side of the neck. Subcutaneous injections should be made under the skin high on the neck behind the ear.</li> </ol>
---------------------	---

Withholding Periods:	<p>MEAT: Zero (0) days  MILK: Zero (0) days</p>
----------------------	---

Trade Advice:	<p>TRADE ADVICE—EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days. Before using this product, confirm the current ESI from Elanco Australasia Pty Ltd on 1800 995 709 or the APVMA website (<a href="http://www.apvma.gov.au/residues">www.apvma.gov.au/residues</a>)</p>
---------------	--

Safety Directions:	
--------------------	--

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26</p>
-------------------------	--

First Aid Warnings:	
---------------------	--

Additional User Safety:	
-------------------------	--

Environmental Statements:	
---------------------------	--

Disposal:	<p>Dispose of empty container by wrapping with paper and putting in garbage.</p>
-----------	--

Storage:

Store below 30°C (Room Temperature). Protect from light. Do not freeze.  
Broached products should be stored below 25°C (Air Conditioning).  
Dispose of broached vials within 28 days of opening.