

READ LABEL BEFORE USE

Combustible liquid. May be fatal if swallowed. May be harmful if absorbed through the skin. May be harmful if inhaled. Causes serious eye irritation and mild skin irritation. Use personal protective equipment as required. If medical advice is needed, have product container or label at hand. Propetamphos is a cholinesterase inhibitor and causes neurotoxicity. Symptoms of poisoning include excessive sweating, nausea, giddiness, headache, weakness, vomiting, contracted pupils, blurred vision, and salivation.

HANDLING PRECAUTIONS

Keep away from sources of ignition (heat, sparks and open flames). Avoid contact with skin or eyes and inhalation of spray mist. When handling concentrate wear PVC or NEOPRENE gloves, waterproof clothing and face shield. Wash splashes of concentrate from skin immediately. Wash protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

If swallowed do NOT induce vomiting but immediately call the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If inhaled, move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped. Use mouth-to nose rather than mouth-to mouth. Obtain medical attention. If in eyes rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye or skin irritation persists or if you feel unwell get medical attention. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately

ENVIRONMENTAL PROTECTION

Very toxic to aquatic organisms with long lasting effects. Toxic to terrestrial vertebrates. Very toxic to terrestrial invertebrates. Avoid contamination of any water supply with product or empty container. Avoid release to the environment, Collect spillage.

EMERGENCY RESPONSE

In case of spillage wear appropriate protective clothing and prevent material from entering water ways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely. For specialist advice in an emergency call **0800 734 607**

This product must be under the personal control of an approved handler or



5 LITRES

PRODUCT INFO

BLUE #:	700749APB	Product Name:	SERAPHOS 1250
Item Code:	YL700749B	Component:	Front Label
Product Code:	N/A	Pack Size:	5L
Provious Itom Code	8/168/75		

ARTWURK INF	U							
Template:		265mm (W) x 175mm (H	Packag	jing Spec(s)	:		N/A
Barcodes/Type: EAN 13: 94210122912		421012291216	Add. In	Add. Info:			N/A	
	Minimum Core Data Point Size:			N/A				
GTIN:		GT	IN Not Required	Proof #	P1a	By/Date	EC	01-N0V-2021
NON PRINTING	CYAN	MAGENTA	YELLOW	BLACK	PANTONE 10	9 C PANTONE	2935 C	PANTONE 2995 C

GENERAL RECOMMENDATIONS

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

For maximum protection: \bullet Saturate the fleece to the skin \bullet Dip 6 – 8 weeks after shearing. \bullet For accurate dipping, calibrate sump, tanks and sticks in 200L steps.

MIXIN

To obtain a uniform wash, premix the measured amount of dip concentrate in a bucket of water. Add the premix to near empty sump or tank. Fill with water. Mix thoroughly.

WITHHOLDING PERIOD

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Animals producing meat and offal for human consumption must not be sold for slaughter during or within **14 days** of the last treatment.

Milk intended for sale for human consumption must be discarded during treatment and for not less than **35 days** following the last treatment.

WOOL RESIDUES

Ensure the time between dipping and subsequent shearing is at least 60 days except in the case of spot treatment of flystrike. For advice on your dipping programme, contact your local supplier.

CONSTANT REPLENISHMENT SHOWER

- Fill sump and tanks and circulate through nozzles to complete mixing.
- FOR BEST RESULTS EMPTY, CLEAN AND REFILL THE SUMP AFTER DIPPING 1 SHEEP PER 2 LITRES OF INITIAL SUMP VOLUME.
 E.G. EMPTY 1,000L SUMP AFTER DIPPING 500 SHEEP.
- Keep sump full at all times.
- Clean out sump after each day's dipping.
- Constant replenishment does not overcome the sump contamination problem.

	Fly & Cattle Tick control	Fly & Cattle Tick contr
d supply tank	1:2,000 or	1:1,250 or

Lice, Ked & LIMITED Lice, Ked & MAXIMUM

MINISHOWER, SPRAY RACE etc

Supply tank	1:800 or 125 mL/100 L water
-------------	-----------------------------

Warning

Sump and

This method of dipping will provide a shorter protection period compared with CR shower or plunge dipping.

YL700749C

JETTING

 For flystrike control, saturate using a handpiece specifically designed for jetting (not a gorse gun).
 DILUTION RATE: 1:800 or 125 mL/100 L water.

STOCK MANAGEMENT

BATCH:

EXPIRY:

- To avoid BLOOD POISONING wait 10 days after shearing for cuts to heal.
 Remove sharp projections from dip area. Do not use biting dogs.
 Vaccinate at least 2 weeks before dipping to protect against Blackleg,
 Malignant Oedema and Pulpy Kidney.
- To avoid PNEUMONIA do not dip sheep that are tired, heated or thirsty.
 Dip early in the day so sheep are dry when returned to pasture. Wet sheep should not be held on grated floors, in wool sheds or covered yards.
- Muster the day before dipping, withholding feed but NOT water 12 – 24 hours before dipping.
- Separate ewes and lambs before dipping.

PLUNGE DIPS AND STANDARD SHOWERS

	Lice, Ked & LIMITED Fly & Cattle Tick control	Lice, Ked & MAXIMUM Fly & Cattle Tick contro	
Initial charge	1:2,000 or 50 mL/100 L water	1:1,250 or 80 mL/100 L water	
Replenishment	1:1,333 or 75 mL/100 L water	1:800 or 125 mL/100 L water	
Dipping Out	1:2,000 or 50 mL/100 L water	1:1,250 or 80 mL/100 L water	

FOR BEST RESULTS EMPTY, CLEAN AND REFILL THE SUMP AFTER DIPPING 1 SHEEP PER 2L OF INITIAL SUMP VOLUME. E.G. EMPTY 1,000L SUMP AFTER DIPPING 500 SHEEP.

Keep each sheep 15 seconds in the dip and submerge head twice for complete coverage. Replenish with water and concentrate each time the wash level falls by 20% of initial volume.



DIPPING OUT

Never dip out from sumps smaller than 10,000 L. Never dip out to below 50% of initial charge volume, Never dip out if more than 1 sheep per 2 L of initial sump volume has been dipped.

BACTERIOSTAT

To prevent post dipping lameness caused by Erysipelothrix rhusiopathiae. If wash is to be used the next day add a bacteriostat immediately after the day's dipping. BUT only if fewer than 1 sheep per 2 L of sump volume has been dipped.

STORAGE

Store locked up in a well-ventilated place. Keep cool.

DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill/facility. Triple rinse empty container and add rinsate to dipwash. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at an approved landfill/facility. Do not use container for any other purpose.



Registered pursuant to the ACVM Act 1997, No.A4265 See www.foodsafety.govt.nz for registration conditions. Elanco New Zealand

106 Wiri Station Road, Manukau, Auckland 2104. Customer Info Line: 0800 446 121

Seraphos 1250 is the registered trade mark of Nippon Kayaku Co. Ltd, Japan

Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2021 Elanco or its affiliates



TRANSPORT UN3018 ORGANOPHOSPHOROUS PESTICIDE, LIQUID, TOXIC, NOS (36% PROPETAMPHOS)





75mm deep

PRODUCT INFO

BLUE #:	700749APB	Product Name:	SERAPHOS 1250
Item Code:	YL700749C	Component:	Back Label
Product Code:	N/A	Pack Size:	5L
Previous Item Code	86662884		

ARTWORK INFO



ELANCO ARTWORK LEGEND v13

265mm wide