

Product name: Flubenol 5 %
Revision date: 21 Mar 2024

Page: 1 of 12
Country: New Zealand

Text above this line is not included in the actual label.

BOTTLE booklet – Front and Base Panels

WARNING
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

Elanco AF0664
FLUBENOL 5%
Oral wormer for pigs and poultry

ACTIVE INGREDIENT
50 mg/g FLUBENDAZOLE

[Pig and chicken pictograms]

Net Content: 600 g

Meat WHP: Pigs: 7 days; Poultry: NIL

Eggs WHP: Poultry: NIL

STORAGE: Store at 15 – 30 °C (room temperature). Tightly reseal open packs before storage.

Read the enclosed leaflet for full instructions

Registered pursuant to the ACVM Act 1997, No. A006818
See www.foodsafety.govt.nz for registration conditions.

Elanco Australasia Pty Ltd
27/88 Shortland St
Auckland, 1010
www.elanco.co.nz
Phone: 0800 446 121



Batch No.
Expiry

Text above this line is not included in the actual label.

Multi-fold Leaflet (inner)

FOR ANIMAL TREATMENT ONLY

Elanco **FLUBENOL 5%** **Oral wormer for pigs and poultry**

ACTIVE INGREDIENT: 50 mg/g FLUBENDAZOLE

Flubenol 5% is indicated for worm infestations in pigs and poultry.

Pigs: infestations with gastrointestinal worms (*Ascaris suum*, *Hyostromylus rubidus*, *Oesophagostomum dentatum*, *Trichuris suis*, *Strongyloides ransomi*) and lungworms (*Metastrongylus apri*).

Poultry: infestations with gastrointestinal worms (*Ascaridia galli*, *Heterakis gallinarum*, *Capillaria* spp., *Amidostomum anseris* and *Trichostrongylus tenuis*, *Raillietina* sp.) and gapeworm (*Syngamus trachea*).

DOSAGE

PIGS

For treatment of piglets and fatteners

30 g flubendazole per ton of feed (30 ppm) for 5 consecutive days. In cases of severe infestation with *T. suis*, the animals should be treated for 10 days.

For treatment of breeding pigs

30 g flubendazole per ton of feed (30 ppm) for 10 consecutive days.

Individual and single administration to breeding pigs

Flubenol 5% is strewn over the feed in a dose of 5 mg per kg of body weight. This corresponds to 1 measuring spoon per 130 kg body weight.



Text above this line is not included in the actual label.

POULTRY

Turkey

20 g flubendazole per ton of feed (20 ppm) for 7 consecutive days.

Chickens and geese

30 g flubendazole per ton of feed (30 ppm) for 7 consecutive days.

In case of infestation with tapeworm (*Rallietina* sp.) give 60 g flubendazole per ton of feed (60 ppm) for 7 consecutive days.

Pheasants and partridges

60 g flubendazole per ton of feed (60 ppm) for 7 consecutive days.



A handwritten signature in blue ink, appearing to read "Barayan".

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Flubenol 5% should be mixed into the feed and the mixture administered.

For a group treatment, make sure that all animals take in enough Flubenol 5%. The medicine should therefore be well mixed into the feed so that a homogenous mixture is obtained. It should not be strewn over pellets or crumbs since this gives a nonhomogeneous mixture.

- 20 g flubendazole per ton (20 ppm) corresponds to 400 g of Flubenol 5% per ton of feed.
- 30 g flubendazole per ton (30 ppm) corresponds to 600 g of Flubenol 5% per ton of feed.
- 60 g flubendazole per ton (60 ppm) corresponds to 1.2 kg of Flubenol 5% per ton of feed.

For small numbers of animals, 1 measuring spoon of Flubenol 5% is mixed into the following amounts of compound feed for the following species: pheasants and partridges: 11 kg; chickens and geese: 21.5 kg; turkeys: 32 kg.

Important: Treatment with Flubenol 5% gives optimal results only when strict hygiene of the pig sheds and the poultry houses is practiced.

Text above this line is not included in the actual label.

CONTRAINDICATIONS

The use of Flubenol 5% is contraindicated in pigeons and parrots.

SIDE EFFECTS

In pigs an overdose in the feed of 250 ppm flubendazole or higher (*i.e.* more than eight times the normal dose) can bring about short-lasting diarrhea (soft or watery dung), without other clinical signs or effects with reference to the growth of the animals. This loose dung occurs from the second treatment day and is maximal after around 7 – 12 days.

In poultry even three times the highest dose (180 ppm flubendazole) for 7 days had no negative effects on egg production, egg quality, hatching results, growth of offspring or in any other respect.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS

PIGS

MEAT: Pigs producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 7 days of the last treatment.

POULTRY (CHICKENS, GAME BIRDS, TURKEYS AND GEESE)

MEAT: NIL

EGGS: NIL



Text above this line is not included in the actual label.

EPA statements

READ ENTIRE LABEL BEFORE USE

WARNING

May cause damage to organs through prolonged or repeated exposure.

HANDLING PRECAUTIONS

Do not breathe dust. May form combustible dust concentrations in air.



FIRST AID

Get medical advice/attention if you feel unwell. For advice contact the National Poisons Centre 0800 POISON (0800 764 766). If medical advice is needed, have the product container or label at hand.

EMERGENCY RESPONSE

Avoid dispersal of spilt material and runoff and contact with soil, waterways, drains and sewers. Use spark-proof tools and explosion-proof equipment. Avoid dust generation. Do not dry sweep. For specialist advice in an emergency call CHEMTREC 0800 425 459. If additional hazard information is required, refer to the Safety Data Sheet. For a copy please go to www.elanco.co.nz.

ENVIRONMENTAL PROTECTION

Avoid release to the environment. Collect spillage.

DISPOSAL

Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Dispose in accordance with local requirements.

STORAGE: Flubenol 5% should be kept at 15 – 30 °C (room temperature).

Tightly reseal open packs before storage.

Feed prepared with Flubenol 5% can be kept for up to 8 weeks.

EXCLUSION OF LIABILITY

To the extent permitted by law, Elanco Australasia Pty Ltd will not be liable or responsible for any loss, damage or expense where the product is not used as directed.

Registered pursuant to the ACVM Act 1997, No. A006818

See www.foodsafety.govt.nz for registration conditions.

Manufactured in the UK.



Product name: Flubenol 5 %
Revision date: 21 Mar 2024

Page: 6 of 12
Country: New Zealand

Text above this line is not included in the actual label.

Registered to
Elanco Australasia Pty Ltd
27/88 Shortland St
Auckland, 1010

Phone: 0800 446 121
Net Content: 600 g

[Trademark statement]



Product name: Flubenol 5 %
Revision date: 21 Mar 2024

Page: 7 of 12
Country: New Zealand

Text above this line is not included in the actual label.

BAG – Front Panel

WARNING
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

Elanco
FLUBENOL 5%

AF0664

Oral wormer for pigs and poultry

ACTIVE INGREDIENT
50 mg/g FLUBENDAZOLE

Batch No.
Expiry

[Pig and chicken pictograms]

Net Content: 2 kg / 25 kg

Registered pursuant to the ACVM Act 1997, No. A006818
See www.foodsafety.govt.nz for registration conditions.

Elanco Australasia Pty Ltd
27/88 Shortland St
Auckland, 1010
Phone: 0800 446 121



A handwritten signature in blue ink, appearing to read "Davyson".

Text above this line is not included in the actual label.

BAG – Back panel

FOR ANIMAL TREATMENT ONLY

Elanco™
FLUBENOL™ 5%
Oral wormer for pigs and poultry

ACTIVE INGREDIENT: 50 mg/g FLUBENDAZOLE

Flubenol 5% is indicated for worm infestations in pigs and poultry.

Pigs: infestations with gastrointestinal worms (*Ascaris suum*, *Hyostrongylus rubidus*, *Oesophagostomum dentatum*, *Trichuris suis*, *Strongyloides ransomi*) and lungworms (*Metastrongylus apri*).

Poultry: infestations with gastrointestinal worms (*Ascaridia galli*, *Heterakis gallinarum*, *Capillaria* spp., *Amidostomum anseris* and *Trichostrongylus tenuis*, *Raillietina* sp.) and gapeworm (*Syngamus trachea*).

DOSAGE

PIGS

For treatment of piglets and fatteners

30 g flubendazole per ton of feed (30 ppm) for 5 consecutive days. In cases of severe infestation with *T. suis*, the animals should be treated for 10 days.

For treatment of breeding pigs

30 g flubendazole per ton of feed (30 ppm) for 10 consecutive days.

Individual and single administration to breeding pigs

Flubenol 5% is strewn over the feed in a dose of 5 mg per kg of body weight. This corresponds to 1 measuring spoon per 130 kg body weight.



Text above this line is not included in the actual label.

POULTRY

Turkey

20 g flubendazole per ton of feed (20 ppm) for 7 consecutive days.

Chickens and geese

30 g flubendazole per ton of feed (30 ppm) for 7 consecutive days.

In case of infestation with tapeworm (*Rallietina* sp.) give 60 g flubendazole per ton of feed (60 ppm) for 7 consecutive days.

Pheasants and partridges

60 g flubendazole per ton of feed (60 ppm) for 7 consecutive days.



DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Flubenol 5% should be mixed into the feed and the mixture administered.

For a group treatment, make sure that all animals take in enough Flubenol 5%. The medicine should therefore be well mixed into the feed so that a homogenous mixture is obtained. It should not be strewn over pellets or crumbs since this gives a nonhomogeneous mixture.

- 20 g flubendazole per ton (20 ppm) corresponds to 400 g of Flubenol 5% per ton of feed.
- 30 g flubendazole per ton (30 ppm) corresponds to 600 g of Flubenol 5% per ton of feed.
- 60 g flubendazole per ton (60 ppm) corresponds to 1.2 kg of Flubenol 5% per ton of feed.

For small numbers of animals, 1 measuring spoon of Flubenol 5% is mixed into the following amounts of compound feed for the following species: pheasants and partridges: 11 kg; chickens and geese: 21.5 kg; turkeys: 32 kg.

Important: Treatment with Flubenol 5% gives optimal results only when strict hygiene of the pig sheds and the poultry houses is practiced.

Text above this line is not included in the actual label.

CONTRAINDICATIONS

The use of Flubenol 5% is contraindicated in pigeons and parrots.

SIDE EFFECTS

In pigs an overdose in the feed of 250 ppm flubendazole or higher (*i.e.* more than eight times the normal dose) can bring about short-lasting diarrhea (soft or watery dung), without other clinical signs or effects with reference to the growth of the animals. This loose dung occurs from the second treatment day and is maximal after around 7 – 12 days.

In poultry even three times the highest dose (180 ppm flubendazole) for 7 days had no negative effects on egg production, egg quality, hatching results, growth of offspring or in any other respect.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS

PIGS

MEAT: Pigs producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 7 days of the last treatment.

POULTRY (CHICKENS, GAME BIRDS, TURKEYS AND GEESE)

MEAT: NIL

EGGS: NIL



Text above this line is not included in the actual label.

EPA statements

READ ENTIRE LABEL BEFORE USE

WARNING

May cause damage to organs through prolonged or repeated exposure.

HANDLING PRECAUTIONS

Do not breathe dust. May form combustible dust concentrations in air.



FIRST AID

Get medical advice/attention if you feel unwell. For advice contact the National Poisons Centre 0800 POISON (0800 764 766). If medical advice is needed, have the product container or label at hand.

EMERGENCY RESPONSE

Avoid dispersal of spilt material and runoff and contact with soil, waterways, drains and sewers. Use spark-proof tools and explosion-proof equipment. Avoid dust generation. Do not dry sweep. For specialist advice in an emergency call CHEMTREC 0800 425 459. If additional hazard information is required, refer to the Safety Data Sheet. For a copy please go to www.elanco.co.nz.

ENVIRONMENTAL PROTECTION

Avoid release to the environment. Collect spillage.

DISPOSAL

Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Dispose in accordance with local requirements.

STORAGE

Flubenol 5% should be kept at 15 – 30 °C (room temperature). Tightly resealed open packs before storage.

Feed prepared with Flubenol 5% can be kept for up to 8 weeks.

EXCLUSION OF LIABILITY

To the extent permitted by law, Elanco Australasia Pty Ltd will not be liable or responsible for any loss, damage or expense where the product is not used as directed.

Registered pursuant to the ACVM Act 1997, No. A006818
See www.foodsafety.govt.nz for registration conditions.



Product name: Flubenol (A006818)
Revision date: 21 Mar 2024

Page: 12 of 12
Country: New Zealand

Text above this line is not included in the actual label.

Manufactured in the UK.

Registered to
Elanco Australasia Pty Ltd
27/88 Shortland St
Auckland, 1010
Phone: 0800 446 121

[Trademark statement]

