Carton front panel	Issue Date: 25 Sept 20	
A single item from each conjoined sets of square brackets i.e. [][] is used per pack.		
The information above this line is not included in the label		

FOR ANIMAL TREATMENT ONLY

DRONTAL® ALLWORMER FOR CATS 6KG

Kills all gastrointestinal worms: Roundworm, Hookworm and Tapeworm (including the Common Flea Tapeworm)

One tablet per 6kg cat bodyweight

Each tablet contains 345mg pyrantel embonate and 30mg praziquantel

[2][4] Tablets

[2][4] tablet packs.....for round shaped tablets
[2] tablet pack.....for ellipsoidal tablets



Carton back panel	Issue Date: 25 Sept 20
The information above this line is not included in the label	

Treatment Programmes

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Kittens: Treat at 4, 6, 8 and 12 weeks of age, then every 3 months or on veterinary advice.

Adult cats: Treat every 3 months. Flea tape worms (*Dipylidium*) can reinfest cats and begin producing segments in faeces 2-3 weeks after treatment. If retreatment is needed before the next DRONTAL treatment is due, treat with DRONCIT® as an intermediate tapeworm-only treatment.

Pregnant Queens: Treat just prior to mating, then 10 days before kittening and at 2 and 4 weeks after kittening. DRONTAL may be given to pregnant queens at any stage of pregnancy.

Note: DRONTAL does not control lungworm.

DOSAGE AND ADMINISTRATION

DRONTAL is given as a single dose by mouth. The tablet is best administered as whole portions rather than crushed in feed since praziquantel gives it a bitter taste. DRONTAL may be given on an empty stomach or with food.

The dose rate is 1 tablet per 6 kg liveweight. Doses should be rounded up to the nearest ½ tablet.

Bodyweight	Tablets
3 kg and under	1/2
3.1-6 kg	1
6.1-9 kg	1 1/2
9.1-12 kg	2

Safety

Drontal may be used concurrently with other treatments, such as Advantage for flea control. Dose rates 5x those recommended have been readily tolerated in clinical studies. In extremely rare cases cats may show slight and transient incoordination. Recovery is always complete within 24 hours and no specific treatment is required.



Carton back panel (cont'd) | Issue Date: 25 Sept 20

The information above this line is not included in the label

Storage

Store below 30°C (Room temperature). Keep out of reach of children.

DRONTAL ALLWORMER FOR CATS 6KG is registered pursuant to the ACVM Act 1997 No. A8215

See www.foodsafety.govt.nz for registration conditions.

DRONTAL, DRONCIT and ADVANTAGE are registered trade marks of Elanco ADVANTAGE ACVM No. A7522, DRONCIT ACVM No. A2919

Elanco New Zealand Ltd 106 Wiri Station Road, Manukau, Auckland 2104

Customer info line: 0800 446 121

Batch No: Expiry:



Cart	on fror	nt pane	el 💮	Issu	e Da	ate: 2	25 Sep	t 20	
		_							П

A single item from each conjoined sets of square brackets i.e. [][] is used per pack.

The information above this line is not included in the label

FOR ANIMAL TREATMENT ONLY

DRONTAL® ALLWORMER FOR CATS 6kg

Tablets for the control of Roundworm, Hookworm and Tapeworm in cats

Each tablet contains 345 mg pyrantel embonate and 30 mg praziquantel

100/ 96 tablets (Elanco logo)

1 tablet per 6 kg

100 tablet pack...... for round shaped tablets 96 tablet pack......for ellipsoidal tablets



Carton back panel	Issue Date: 25 Sept 20	
The information above this line is not included in the label		

TREATMENT PROGRAMMES

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Kittens: Treat at 4, 6, 8 and 12 weeks of age, then every 3 months or on veterinary advice.

Adult cats: Treat every 3 months. Flea tape worms (*Dipylidium*) can reinfest cats and begin producing segments in faeces 2-3 weeks after treatment. If retreatment is needed before the next DRONTAL treatment is due, treat with DRONCIT® as an intermediate tapeworm-only treatment.

Pregnant Queens: Treat just prior to mating, then 10 days before kittening and at 2 and 4 weeks after kittening. DRONTAL may be given to pregnant queens at any stage of pregnancy.

Note: DRONTAL does not control lungworm.

DOSAGE AND ADMINISTRATION

DRONTAL is given as a single dose by mouth. The tablet is best administered as whole portions rather than crushed in feed since praziquantel gives it a bitter taste. DRONTAL may be given on an empty stomach or with food. The dose rate is 1 tablet per 6 kg liveweight. Doses should be rounded up to the nearest ½ tablet.

Bodyweight	Tablets
3 kg and under	1/2
3.1-6 kg	1
6.1-9 kg	1 1/2
0.1.12 kg	2

SAFETY

Drontal may be used concurrently with other treatments, such as Advantage for flea control. Dose rates 5x those recommended have been readily tolerated in clinical studies. In extremely rare cases cats may show slight and transient incoordination. Recovery is always complete within 24 hours and no specific treatment is required.



Carton back panel (cont'd) | Issue Date: 25 Sept 20

The information above this line is not included in the label

STORAGE

Store below 30°C (Room temperature). KEEP OUT OF REACH OF CHILDREN.

DRONTAL ALLWORMER FOR CATS 6KG is registered pursuant to the ACVM Act 1997 No. A8215

See www.foodsafety.govt.nz for registration conditions.

DRONTAL, DRONCIT and ADVANTAGE are registered trade marks of Elanco ADVANTAGE ACVM No. A7522, DRONCIT ACVM No. A2919

Elanco New Zealand 106 Wiri Station Road, Manukau, Auckland 2104

Customer info line: 0800 446 121

Batch No: Expiry:



Package Insert | Issue Date: 25 Sept 20 The information above this line is not included in the label

DRONTAL® ALLWORMER FOR CATS 6KG

FOR ANIMAL TREATMENT ONLY

Kills all gastrointestinal worms: Roundworm, Hookworm and Tapeworm (including the Common Flea Tapeworm)

One tablet per 6kg cat bodyweight

Each tablet contains 345mg pyrantel embonate and 30mg praziquantel

DRONTAL ALLWORMER FOR CATS 6 KG is for the control of all gastrointestinal worms including tapeworms in cats:

Roundworms

Toxocara cati Toxascaris leonina **Hookworms**

Uncinaria stenocephala*

Tapeworms

Dipylidium caninum (common flea tapeworm)

Taenia taeniaeformis

*Not recognised as clinically significant in cats in New Zealand

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

DRONTAL IS GIVEN AS A SINGLE DOSE BY MOUTH.

It is essential to weigh cats before treatment.

Worms, if present, may not appear in the droppings after treatment, or may not appear for up to 48 hours after treatment. DRONTAL is given as a single dose by mouth. The tablet is best administered as whole portions rather than crushed in feed since praziquantel gives it a bitter taste. DRONTAL may be given on an empty stomach or with food.



DOSAGE

Bodyweight	Tablets
3kg and under	1/2
3.1kg – 6kg	1
6.1kg – 9kg	1 1/2
9.1kg – 12kg	2

DOSING TIPS

Cats are dosed according to weight so weigh your cat before each treatment. The dose rate is 1 tablet per 6kg bodyweight. Doses should be rounded up to the nearest ½ tablet.

RECOMMENDED TREATMENTS

Worms often reinfest cats so it is necessary to treat the cat itself and reduce environmental contamination by daily disposal of droppings where possible. Also control intermediate hosts, eg. fleas, rats and mice. This table summaries treatment recommendations:

KITTENS	ADULTS	Pregnant/Lactating
		Queens
Treat at 4, 6, 8, 12 weeks; then 3 monthly	Every 3 months	Prior to mating, 10 days before kittening and 2 and 4 weeks after kittening. Then every 3
		months.

These recommendations are of a general nature. In specific instances where infestation is particularly heavy, or many cats are housed together, a veterinary surgeon should be consulted. Flea tapeworms can reinfect cats and begin producing segments 2-3 weeks after treatment. If segments reappear before the next DRONTAL treatment is due, either retreat with DRONTAL ALLWORMER FOR CATS or use DRONCIT® as an intermediate tapeworm-only treatment.

CONTROL OF REINFESTATION

Roundworm, Hookworm:

Hookworms are usually transmitted to kittens through the mother's milk and it is difficult to prevent this transmission. Both hookworms and roundworms are also transmitted by eggs and larvae on the ground while roundworm can infect cats through rats and mice. The eggs of roundworm are very resistant and may survive in soil for years so regular treatment to prevent environmental contamination with eggs is essential. Kittens under 6 months and pregnant and lactating queens produce excessive quantities of worm eggs and therefore require more attention to worm control.

Tapeworm:

Because these worms infest cats through a carrier host, reinfestation can be delayed or controlled by preventing access to common carrier hosts such as rats, mice and fleas. A good flea control programme will help control flea tapeworm.

COMPATIBILITY

Drontal may be used concurrently with other treatments, such as Advantage for flea control.

SAFETY

DRONTAL offers gentle, effective control of worms. It may be given to all cats including old animals, young kittens, pregnant queens and show animals. In extremely rare cases cats may show slight and transient incoordination. Recovery is always complete within 24 hours and no specific treatment is required.

STORAGE

Store below 30°C (Room temperature). Keep out of reach of children

Drontal ALLWORMER FOR CATS 6KG is registered pursuant to the ACVM Act 1997 No. A8215

See www.foodsafety.govt.nz for registration conditions.

Elanco New Zealand 106 Wiri Station Road, Manukau, Auckland 2104

Customer info line: 0800 446 121

DRONTAL®, DRONCIT® and ADVANTAGE® are trade marks of ElancoDRONCIT ACVM No. A2919 ADVANTAGE ACVM No. A7522

