

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY



Elanco seresto

for Dogs and Puppies

Fleas & ticks



OR



ACTIVE CONSTITUENTS:

100 mg/g IMIDACLOPRID and 45 mg/g FLUMETHRIN.

Treats and prevents flea infestations (*Ctenocephalides felis*) for 8 months. Controls paralysis ticks (*Ixodes holocyclus*) for 8 months.

Repels paralysis ticks (*Ixodes holocyclus*) for up to 4 months. Controls and reduces incidence of flea allergy dermatitis (FAD). Kills larval flea stages in the surroundings of treated dogs. Fleas already on dogs prior to treatment will be killed within 24 hours after collar application. After 7 days of collar application new flea infestations will be killed within 2 hours. Prevents flea tapeworm (*Dipylidium caninum*) infection through effective flea control.

Controls brown dog ticks (*Rhipicephalus sanguineus*) and bush ticks (*Haemaphysalis longicornis*) for 4 months.

Effective against larval, nymph and adult stages of brown dog ticks. Reduces viable egg production by female brown dog ticks.

Reduces the transmission between dogs of diseases carried by the brown dog tick: Canine Monocytic Ehrlichiosis CME (due to *Ehrlichia canis*), Babesiosis (due to *Babesia vogeli*) and Anaplasmosis (due to *Anaplasma platys*) for 4 months.

Treats biting/chewing lice (*Trichodectes canis*) present at the time of treatment.

Ixodes holocyclus ticks do not occur in Western Australia.

In tick season, daily searching for, and removal of, any ticks found is recommended. The Seresto collar contains flumethrin, a synthetic pyrethroid, and imidacloprid, a neonicotinoid. If problems persist, consult a veterinarian.

PRECAUTIONS

PREGNANCY AND LACTATION: The safety of this product has not been established during pregnancy and lactation.

SIDE EFFECTS

Application site reactions such as dermatitis, inflammation, eczema or lesions may occur and in these instances, a temporary collar removal may be recommended until the symptoms have disappeared. Also in rare cases, slight and transient reactions such as depression, change of food intake, salivation, vomiting and diarrhoea might occur initially. In the unlikely event of the animal eating the collar, mild gastrointestinal symptoms (e.g. loose stools) may occur.

DOSAGE AND ADMINISTRATION

For external use only


For puppies from 7 weeks of age

One collar per animal fastened around the neck.

Small dogs and puppies ≤ 8 kg body weight require one SERESTO TICK AND FLEA COLLAR FOR DOGS AND PUPPIES ≤ 8 KG.

Dogs > 8 kg body weight require one SERESTO FOR DOGS > 8 KG FLEAS & TICKS COLLAR.

1. Remove collar from plastic pouch immediately before use
2. Unroll collar and remove plastic connectors remaining on the inside edge of the collar (these can cause mechanical irritation if not removed)
3. Adjust the collar around the dog's neck without over-tightening. As a guide, it should be possible to insert 2 fingers between the collar and the dog's neck.
4. Pull collar through the loop and cut off any excess length extending beyond 2 cm. In puppies, allow some additional length to allow for loosening of the collar as the puppy grows.



Important: Unroll collar and remove plastic connectors remaining on the inside edge of the collar.

Check periodically and adjust fit if necessary, especially when puppies are rapidly growing.

The collar should be worn continuously for the full protection period. If applying ADVOCATE, ADVANTAGE or ADVANTIX to a dog wearing a SERESTO collar, there is no need to remove the Seresto collar. Follow the label instructions.

GENERAL DIRECTIONS

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

ODOURLESS, WATER RESISTANT.

FLEAS

The Seresto collar treats and control fleas for 8 months. Fleas can be difficult to control, because the majority of the flea population in the home is in the form of flea eggs, larvae and pupae, rather than adult fleas on the dog. Ideally, flea treatment for dogs should start early in their life, as soon as they join your family or farm. Lapses in flea control can allow the flea population to establish or rebuild, leading to an infestation that can take months to eliminate. This

is why it's important to continue using flea control all year-round, and on all pets, even during the cooler winter months.

The Seresto collar also kills flea larvae in your dog's surroundings, however in the case of a heavy flea infestation, environmental sprays, vacuuming and washing pet bedding will also help resolve the problem.

Shampooing, frequent exposure to water (e.g. swimming, bathing or rain) may reduce the collar's efficacy against fleas (*Ctenocephalides felis*). If washing your dog, always use a soap-free shampoo.

Flea tapeworm

The flea tapeworm (*Dipylidium caninum*) is transmitted by fleas. Dogs are infected by ingesting adult fleas infected with the immature stages of this tapeworm. Effective flea control with the Seresto collar prevents flea tapeworm infection. To control other tapeworms, treat as directed by your veterinarian.

TICKS

Ticks already on dogs prior to applying the Seresto collar may not be killed immediately after collar application, and may remain attached and visible. Therefore removal of ticks already on the dog at the time of application is recommended. Effective control of new brown dog tick infestations is achieved by 48 hours following collar application.

Shampooing and frequent exposure to water does not reduce the efficacy of Seresto against the brown dog tick (*Rhipicephalus sanguineus*). The effect of shampooing or exposure to water on efficacy against paralysis tick (*Ixodes holocyclus*), bush tick (*Haemaphysalis longicornis*) or lice (*Trichodectes canis*) has not been established.

Paralysis tick

The paralysis tick (*Ixodes holocyclus*) is a potentially deadly parasite and difficult to control. While the Seresto collar repels paralysis ticks for up to 4 months and controls paralysis ticks for 8 months, following collar application the risk of tick paralysis cannot be entirely eliminated as it takes only one adult female tick to cause tick paralysis/poisoning. Therefore, daily searching for and removal of any ticks found is recommended.

VECTOR-BORNE DISEASES

The Seresto collar reduces the transmission of diseases spread by the brown dog tick, including ehrlichiosis, babesiosis and anaplasmosis for 4 months.

PUPPIES

Puppies may be treated with a Seresto collar from 7 weeks of age. If puppies younger than 7 weeks require flea control they may be safely treated with ADVANTAGE for dogs.

COLLAR TECHNOLOGY - POLYMER MATRIX



The patented polymer matrix system of the Seresto collar ensures that both active ingredients are slowly and continuously released from the collar, as needed, ensuring low acaricidal/insecticidal concentrations on the dog throughout the entire efficacy period. The active ingredients spread from where the collar contacts the skin, across the entire skin surface.

The modes of action do not require systemic absorption of active ingredients into the dog, so efficacy is achieved without requiring fleas or ticks to bite and ingest a blood meal.

This collar is designed with a safety-closure mechanism. In the unlikely event of a dog becoming trapped, the dog's own strength is sufficient to widen the collar to allow for a quick release.

SAFETY DIRECTIONS

Harmful if swallowed. Do not open inner pouch until ready to use. Do not allow children to play with the collar. Wash hands after use. Wear gloves if applying a collar to more than one animal per day.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

ADDITIONAL USER SAFETY

As with any veterinary medicinal product, do not allow small children to play with the collar, or to put it into their mouth. People with known sensitivity to the ingredients of the collar should avoid contact with the collar. Additional information can be located on the Safety Data Sheet (SDS).

DISPOSAL

Dispose of used collars and plastic pouch by wrapping with paper and putting in garbage.

STORAGE Store below 30°C (room temperature)

LIMITATION OF LIABILITY

Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Elanco Australasia Proprietary Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

PRESENTATION

Product Name	Contents	APVMA Approval No.
SERESTO FOR DOGS AND PUPPIES ≤ 8 KG FLEAS & TICKS COLLAR	1 collar – 38 cm length	66209/133078
SERESTO FOR DOGS > 8 KG FLEAS & TICKS COLLAR	1 collar – 70 cm length	65875/133077

Elanco Australasia Pty Ltd
Level 3, 7 Eden Park Dr
Macquarie Park NSW 2113
ACN 076 745 198

CUSTOMER INFORMATION
Australia: 1800 995 709 TOLL FREE
www.elanco.com.au

Seresto, Advantage, Advocate, Advantix, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.
© 2021 Elanco or its affiliates.

Elanco