



WARNING
KEEP OUT OF REACH OF CHILDREN
READ LABEL BEFORE USE
FOR ANIMAL TREATMENT ONLY

Elanco Seresto™

8 Month Flea and Tick Collar for Cats and Kittens
 38 cm – 12.5 g

Active Constituents:

Each collar contains: imidacloprid 100 mg/g and flumethrin 45 mg/g.

The patented Bayer polymer matrix system ensures that both active ingredients are slowly and continuously released in low concentrations from the collar onto the animal. This avoids peak concentrations and ensures that the active ingredients are present on the pet's skin and hair for the full duration of the efficacy period.

Fleas:

- For the treatment and prevention of flea (*Ctenocephalides* spp.) infestation for 8 months.
- Efficacy against fleas starts immediately after application of the collar.
- Protects treated pets immediate surroundings against flea larvae development.
- For treatment and prevention of flea allergy dermatitis (FAD).

Ticks:

- Controls ticks (*Haemaphysalis longicornis*) by killing and repelling them for 8 months.
- It is effective against larvae, nymphs and adult tick stages.
- Ticks already on the pet prior to treatment may not be killed immediately after collar application and may remain attached and visible. Therefore removal of ticks already on pets at the time of application is recommended. The prevention of infestations with new ticks starts immediately after application of the collar.
- Ticks will be killed and fall off the host within 24 to 48 hours after infestation without having a blood meal, as a rule. An attachment of single ticks after treatment cannot be excluded.
- As in long term topical products, periods of excessive seasonal hair shedding may lead to transient slight reduction of efficacy by loss of hair-bound portions of the active ingredients. Replenishment from the collar starts immediately so that full efficacy will be re-established without any additional treatment or collar replacement.

Also for treatment of lice infestation in dogs for 8 months.

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain or distress when using this product other than as directed on the label.

DOSAGE

One collar per animal to be fastened around the neck.

[Puppies and small dogs up to 8 kg body weight receive one collar of 38 cm length.

Dogs above 8 kg receive one Seresto collar for dogs over 8 kg of 70 cm length.] [Cats, puppies and small dogs up to 8 kg body weight receive one collar of 38 cm length.

Dogs above 8 kg receive one Seresto collar for dogs over 8 kg of 70 cm length.]

GENERAL INSTRUCTIONS

Remove collar from protective bag directly before use. Unroll collar and make sure that there are no remnants from the plastic connectors inside the collar. Adjust the collar around the animal's neck without tightening it too tight (as a guide, it should be possible to insert 2 fingers between the collar and the neck). Pull excess collar through the loop and cut off any excess length extending beyond 2 cm.

After application of the collar three reflector clips may be fixed permanently to the collar to increase the animal's visibility in the dark. Take the clips out of the bag. The clips should be evenly distributed on the non-overlapping part of the collar. The clips are correctly applied when a clicking sound is heard. For safety reasons once the reflectors are fixed to the collar the clasp closes permanently and cannot be re-opened. The reflector clips contained herein are only meant to be used in combination with the specific collar and have been tested only for such use and purpose.



The collar should be worn continuously for the 8 month protection period. Check periodically and adjust fit if necessary, especially when [kittens][puppies][kittens or puppies] are rapidly growing. [This collar is designed with a safety-closure mechanism. In the unlikely event of a cat being trapped, the cats own strength is sufficient to widen the collar to allow for quick release.]

The product is water resistant and does not need to be removed during bathing. However, prolonged or repeated exposure of the animal to water or excessive washing should be avoided as the duration of activity may be reduced.

PREGNANCY AND LACTATION

The safety of this product has not been established during pregnancy and lactation. Laboratory studies with either flumethrin or imidacloprid in rats and rabbits have not produced any effects on fertility or reproduction nor shown any teratogenic or foetotoxic effects.

PUPPIES/KITTENS

{Do not use in [puppies less than 7 weeks of age][kittens less than 10 weeks of age]. If [puppies][kittens] younger than [7][10] weeks require flea control they may be treated with ADVANTAGE.} {Do not use in puppies less than 7 weeks of age or kittens less than 10 weeks of age. If puppies younger than 7 weeks or kittens less than 10 weeks require flea control they may be treated with ADVANTAGE.}

PRECAUTIONS

Ensure that the collar is not fitted too tightly. Occasionally mild behavioural disorders that may include scratching at the application site may be observed in animals that are not used to wearing collars on the first few days after fitting. Mild application site reactions such as pruritus, erythema and hair loss may occur uncommonly in cats. Animals usually recover within 1 to 2 weeks without the need for collar removal. In single cases, a temporary collar removal may be recommended until the symptoms have disappeared.

In rare cases in cats, application site reactions such as dermatitis, inflammation, eczema or lesions may occur. In these instances, collar removal is recommended.

In rare cases during the first few days after fitting the collar, mild and transient behavioural signs may occur, e.g. depression or change in appetite which may be associated with gastrointestinal symptoms such as salivation, vomiting and diarrhoea.

Do not use with other insecticides.

The supplied reflector clips are only meant to be used in combination with this specific collar and have been tested only for such use and purpose.

PRESENTATION

One collar per pack

For Cats and Kittens – 38 cm length collar (12.5 g)

CUSTOMER INFORMATION LINE

0800 446 121 TOLL FREE from anywhere in New Zealand
 8.30am to 4.30pm Monday to Friday
animalhealth_nz@elanco.co.nz

STORAGE

Store below 30°C (Room temperature)

DISPOSAL

The product should not enter water courses as it may be dangerous for fish and other aquatic organisms. Any unused product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. Dispose of used collar by wrapping in paper and placing in garbage. ACVM Registration No. A10750

See www.foodsafety.govt.nz for registration conditions.

Elanco New Zealand

106 Wiri Station Road,
 Manukau, Auckland 2104.

Customer Information Line: 0800 446 121.

SERESTO™ and ADVANTAGE™ are registered trademarks of Elanco.

Harmful if swallowed. Prolonged exposure by ingestion could cause damage to skin and the liver. Do not breathe dust from collar. If medical advice is needed, have product container or label at hand. People with known sensitivity to the ingredients of the collar should avoid contact with the collar.

The packaging contains small parts (reflectors) which bear a choking hazard. As with any veterinary medicine, do not allow small children to play with the collar or reflectors, or put them into their mouths.

Handling Precautions Wash hands with cold water thoroughly after fitting the collar. Do not eat, drink or smoke when using this product.

First Aid If swallowed or if you feel unwell get medical advice. Call 0800 POISON (0800 764 766) or a doctor.

Environmental Protection

Very toxic to aquatic life, including fish and other aquatic organisms, with long lasting effects and toxic to the soil environment. Avoid release to the environment, particularly water courses.